

Inhomogeneous Restricted Lattice Walks

Manfred Buchacher¹ and Manuel Kauers^{1*}

¹*Institute for Algebra, Johannes Kepler University, Linz, Austria*

Abstract. We consider inhomogeneous lattice walk models in a half-space and in the quarter plane. For the models in a half-space, we show by a generalization of the kernel method to linear systems of functional equations that their generating functions are always algebraic. For the models in the quarter plane, we have carried out an experimental classification of all models with small steps. We discovered many (apparently) D-finite cases for most of which we have no explanation yet.

Keywords: Functional Equations, Kernel Method, Formal Power Series, D-finiteness

1 Introduction

Given a specific counting problem, it is often easy to write down a functional equation for the corresponding generating function, but it can be quite hard to derive from it something interesting about its solution. Counting problems for restricted lattice walks are a source of functional equations which are neither trivial nor hopeless, and therefore interesting. Using the kernel method, Banderier and Flajolet [2] have shown that the generating functions for lattice walks restricted to a half-space are always algebraic. For walks restricted to the quarter plane, the functional equations are more intricate, and as a consequence, the resulting generating functions come in more flavours. A systematic classification was initiated by Bousquet-Mélou and Mishna [12] and has since led to a substantial amount of literature, see [8] and the references given there, as well as [4, 16] for some more recent developments.

Since the main questions raised in the paper of Bousquet-Mélou and Mishna have been answered, the focus has shifted to modified versions of the problem, including, for example, weighted steps [18, 14], longer steps [9], higher dimensions [10, 1, 5], or interacting boundaries [3]. All these variations are homogeneous in the sense that the walks are formed from a fixed set of admissible steps. Little is known about inhomogeneous models, e.g., lattice walk models where the set of admissible steps may vary with the time and/or the position of the walk. Bousquet-Mélou and Xin [13] have studied such a case. They analyzed a quarter plane model where the step set varies with the parity of the time. Also D'Arco et al. [15] have studied an inhomogeneous case. They determined the asymptotics for walks in the quarter plane where at position (i, j) , the next step is

*Both authors were supported by the Austrian FWF grant F5004.

taken from \leftrightarrow if $i + j$ is even and from \bowtie if $i + j$ is odd. In the present paper, we study such inhomogeneous models more systematically.

We consider half-space models and quarter plane models. For the half-space, we show for a general class of inhomogeneities that the generating function is invariably algebraic. Our proof is an adaption of an argument given by Bousquet-Melóu and Jehanne [11] to the case of linear systems of functional equations (Section 3). The statement also follows from a deep theorem of Popescu [19] (see also Theorem 16 in [7]). For regions defined by more than one restriction, we generalize the notion of dimension introduced by Bostan et al. [10] to inhomogeneous models (Section 4). For the quarter plane (Section 5), we give an experimental classification for models with short steps and two specific inhomogeneities. We recognize many generating functions as D-finite, but in most cases we have no explanation for their D-finiteness.

2 Inhomogeneous Lattice Walks in the Half-Space

We consider walks in a half-space $\mathbb{Z}_{\geq 0} \times \mathbb{Z}^{d-1}$ which start at some point (i_0, \mathbf{j}_0) of the half-space, consist of n steps, and end at some point (i, \mathbf{j}) of the half-space. Restrictions are imposed on the steps the walks consist of. For a *homogeneous* model, there is a fixed finite set $\mathbf{S} \subseteq \mathbb{Z}^d$, called the step set, from which each step of the walk is taken.

For an *inhomogeneous* model, the allowed steps are governed by a deterministic finite automaton. A finite automaton is a directed multigraph $(\mathcal{Q}, \mathcal{E})$ whose (finitely many) edges are labelled by letters of some alphabet. The vertices $q \in \mathcal{Q}$ are called states. One particular state $q_0 \in \mathcal{Q}$ is called the initial state, and there is a subset $\bar{\mathcal{Q}} \subseteq \mathcal{Q}$ of final states. In our setting the edges are labelled by elements of \mathbb{Z}^d . To be deterministic means that for every pair (q, \mathbf{s}) with $q \in \mathcal{Q}$ and $\mathbf{s} \in \mathbb{Z}^d$ there is at most one edge starting from q and labelled by \mathbf{s} . A lattice walk $w = w_0, \dots, w_n$ belongs to the inhomogeneous model if $w_k \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}^{d-1}$ for $k = 0, \dots, n$ and there is a path in the automaton starting at the initial state, ending at one of the final states, and such that the k th edge of the path is labelled with $w_k - w_{k-1} \in \mathbb{Z}^d$ for $k = 1, \dots, n$. Below we will write \mathbf{S}_{pq} for the set of all $s \in \mathbb{Z}^d$ which label an edge from p to q .

Example 1. Let $\mathbf{S}_0 = \leftrightarrow$ and $\mathbf{S}_1 = \bowtie$, and consider walks in $\mathbb{Z}_{\geq 0} \times \mathbb{Z}$ starting at $(0, 0)$ and consisting of steps taken from $\mathbf{S}_0 \cup \mathbf{S}_1$ with the restriction that whenever the current position of a walk is (i, j) , the next step must be taken from $\mathbf{S}_{(i+j) \bmod 2}$. In the notation from above we can take $\mathcal{Q} = \bar{\mathcal{Q}} = \{0, 1\}$, $q_0 = 0$, and $\mathbf{S}_{01} = \leftrightarrow$, $\mathbf{S}_{10} = \leftrightarrow$ and $\mathbf{S}_{11} = \bowtie$.

When the admissible steps only depend on the current position (i, \mathbf{j}) , as in Example 1, we call the model *space inhomogeneous*. When they only depend on the time n , the model is called *time inhomogeneous*.

Fix an inhomogeneous lattice walk model as introduced above, and for each set \mathbf{S}_{pq} fix a function $w_{pq} : \mathbf{S}_{pq} \rightarrow \mathbb{K}$, where \mathbb{K} is a field of characteristic zero. We call $w_{pq}(u, \mathbf{v})$

the *weight* of the step $(u, \mathbf{v}) \in \mathbf{S}_{pq}$. The weight of a walk is the product of the weights of the steps it consists of. Note that the weight does not only depend on the step (u, \mathbf{v}) but also on the edge it labels.

For any i, \mathbf{j}, n , let $f_{i, \mathbf{j}, n}$ be the sum of the weights of all walks from (i_0, \mathbf{j}_0) to (i, \mathbf{j}) of length n , and let $F(x, \mathbf{y}, t) = \sum_{i, \mathbf{j}, n} f_{i, \mathbf{j}, n} x^i \mathbf{y}^{\mathbf{j}} t^n \in \mathbb{K}[x, \mathbf{y}, \mathbf{y}^{-1}][[t]]$ be the corresponding generating function, where we write \mathbf{y} and \mathbf{y}^{-1} for y_1, \dots, y_{d-1} and $y_1^{-1}, \dots, y_{d-1}^{-1}$, respectively, and $\mathbf{y}^{\mathbf{j}}$ for $y_1^{j_1} \cdots y_{d-1}^{j_{d-1}}$. Note that $f_{i, \mathbf{j}, n}$ is just the total number of walks in the model if all weights are defined as 1. If F_q is the generating function of walks associated with paths in the automaton ending at state q , then $F = \sum_{q \in \mathcal{Q}} F_q$.

Example 2. Continuing the previous example, F_0 is the generating function counting walks ending at a point (i, j) with $i + j$ even and F_1 is the generating function counting walks ending at a point (i, j) with $i + j$ odd. The full generating function is $F = F_0 + F_1$.

For the inhomogeneities under consideration the F_q 's satisfy a system of functional equations: writing $S_{pq} := \sum_{(u, \mathbf{v}) \in \mathbf{S}_{pq}} w_{pq}(u, \mathbf{v}) x^u \mathbf{y}^{\mathbf{v}}$, the combinatorial specification, including the boundary condition, translates into the system of functional equations

$$\forall q : F_q = [q = q_0] + t \sum_p ([x^{\geq 0}] S_{pq}) F_p + t \sum_{k \geq 1} \sum_p x^{-k} ([x^{-k}] S_{pq}) [x^{\geq k}] F_p,$$

where $[x^{\geq 0}] S_{pq} \in \mathbb{K}[x, \mathbf{y}, \mathbf{y}^{-1}]$ is the Laurent polynomial obtained from S_{pq} by discarding all terms with negative exponents in x , where $[x^{-k}] S_{pq} \in \mathbb{K}[\mathbf{y}, \mathbf{y}^{-1}]$ is the coefficient of x^{-k} of $S_{pq} \in \mathbb{K}[x, x^{-1}, \mathbf{y}, \mathbf{y}^{-1}]$, and where $[x^{\geq k}] F_p$ denotes the series obtained from F_p by discarding all terms $x^i \mathbf{y}^{\mathbf{j}} t^n$ with $i < k$. The functional equation can also be expressed using the Δ operator defined by $\Delta F(x, \mathbf{y}, t) := (F(x, \mathbf{y}, t) - F(0, \mathbf{y}, t)) / x$. Note that $[x^{\geq k}] F = x^k \Delta^k F$ for all $k \in \mathbb{N}$, so $x^{-k} ([x^{-k}] S_{pq}) [x^{\geq k}] F_p$ simplifies to $([x^{-k}] S_{pq}) \Delta^k F_p$.

Example 3. Continuing the previous example and taking all weights to be 1, we have

$$\begin{aligned} F_0(x, y) &= 1 + t(y + y^{-1} + x)F_1(x, y) + tx^{-1}(F_1(x, y) - F_1(0, y)), \\ F_1(x, y) &= t(y + y^{-1} + x)F_0(x, y) + tx(y + y^{-1})F_1(x, y) \\ &\quad + tx^{-1}(F_0(x, y) - F_0(0, y)) + tx^{-1}(y + y^{-1})(F_1(x, y) - F_1(0, y)). \end{aligned} \quad (2.1)$$

Eliminating $F_0(x, y)$ from these equations leads to the equation

$$\begin{aligned} &(x^2 y^2 - t^2(x + y)^2(1 + xy)^2 - tx(1 + x^2)y(1 + y^2))F_1(x, y) \\ &= txy(x + y)(1 + xy) - txy^2 F_0(0, y) - ty(x(1 + y^2) + t(x + y)(1 + xy))F_1(0, y). \end{aligned} \quad (2.2)$$

which we can solve using the kernel method: the polynomial $x^2 y^2 - t^2(x + y)^2(1 + xy)^2 - tx(1 + x^2)y(1 + y^2) \in \mathbb{Q}(y)[[t]][x]$ has two roots in $\mathbb{Q}(y)[[t]]$, and if we substitute them for x in equation (2.2), we get a system of two linear equations for the two unknown series $F_0(0, y)$

and $F_1(0, \mathbf{y})$. This system turns out to have a unique solution, which implies that $F_0(0, \mathbf{y})$ and $F_1(0, \mathbf{y})$ are algebraic. Consequently, by equation (2.2), also $F_1(x, \mathbf{y})$ is algebraic. Consequently, by equation (2.1), also $F_0(x, \mathbf{y})$ is algebraic. Finally, it follows by algebraic closure properties that $F(x, \mathbf{y}) = F_0(x, \mathbf{y}) + F_1(x, \mathbf{y})$ is algebraic.

For any particular choice of inhomogeneity we can write down an explicit system of functional equations for the auxiliary series F_q , which we can attempt to solve as illustrated in the example above. Potentially, such an attempt could fail, for example because there are too few roots for applying the kernel method, or because there are too many solutions of the linear system for the evaluated auxiliary series $F_q(0, \mathbf{y})$. The next theorem says that we do not need to worry about such problems.

Theorem 1. *The generating function $F(x, \mathbf{y}, t) \in \mathbb{K}[x, \mathbf{y}, \mathbf{y}^{-1}][[t]]$ for a model of inhomogeneous lattice walks and a choice of weight functions w_{pq} as specified above is algebraic over $\mathbb{K}[x, \mathbf{y}, t]$.*

Proof. As argued above, for every q there is a functional equation $F_q = \dots$, where the right hand side is a $\mathbb{K}(\mathbf{y})[x, t]$ -linear combination of 1 and the series $\Delta^i F_q$. These equations together form a system of functional equations which can be written as

$$\mathbf{f} = \mathbf{a} + t \sum_{i=0}^k \mathbf{B}_i \Delta^i \mathbf{f},$$

where \mathbf{f} is the vector of the F_q 's, where \mathbf{a} is a certain explicit vector, and where the \mathbf{B}_i 's are certain explicit matrices with entries in $\mathbb{K}(\mathbf{y})[x, t]$. According to Theorem 2 shown in the next section (applied with $\mathbb{K}(\mathbf{y})$ in place of \mathbb{K}), the unique solution vector of such a system always has algebraic components. This means that every F_q is algebraic, and consequently, the finite sum $F = \sum_{q \in \mathcal{Q}} F_q$ is algebraic too. \square

3 Systems of Linear Functional Equations

The purpose of the present section is to prove the following theorem, which says that the solutions of certain systems of linear functional equations are always algebraic. For a more general statement on systems of polynomial equations see Popescu [19].

Throughout this section, let \mathbb{K} be a field of characteristic zero. Recall from the previous section that $\Delta: \mathbb{K}[x][[t]] \rightarrow \mathbb{K}[x][[t]]$ is defined by $\Delta f(x, t) = \frac{1}{x}(f(x, t) - f(0, t))$. Applied to a vector of series, the action of Δ is meant componentwise.

Theorem 2. *Let $\mathbf{a} \in \mathbb{K}[x][t]^n$ and $\mathbf{B}_i \in \mathbb{K}[x][t]^{n \times n}$ ($i = 0, \dots, k$). Then the functional equation*

$$\mathbf{f} = \mathbf{a} + t \sum_{i=0}^k \mathbf{B}_i \Delta^i \mathbf{f} \tag{3.1}$$

has a unique solution \mathbf{f} in $\mathbb{K}[x][[t]]^n$, and its components are algebraic over $\mathbb{K}[x, t]$.

It is clear that the functional equation has a unique solution in $\mathbb{K}[x][[t]]^n$, because we can compute its coefficients recursively via

$$[t^0]\mathbf{f} = [t^0]\mathbf{a} \quad \text{and} \quad [t^{n+1}]\mathbf{f} = [t^{n+1}]\mathbf{a} + \sum_{i=0}^k \sum_{j=0}^n ([t^j]\mathbf{B}_i) \Delta^i [t^{n-j}]\mathbf{f} \quad (n \in \mathbb{N}).$$

The nontrivial part of the theorem is that the components of the solution are algebraic. For proving this part of the theorem, we may assume without loss of generality that \mathbb{K} is algebraically closed, and we will do so from now on.

Our proof is an adaption of the proof idea of Thm. 3 in [11] to linear systems. We first bring the unevaluated terms $\mathbf{f}(x, t)$ hidden in the delta terms to the left hand side, so that the right hand side only contains $\mathbf{f}(0, t)$ or other evaluated versions of \mathbf{f} . This can be done by translating the delta terms into evaluations of partial derivatives. Using $[x^j]\Delta^i \mathbf{f}(x, t) = [x^{i+j}]\mathbf{f}(x, t)$ and $[x^j]\mathbf{f}(x, t) = \frac{1}{j!} \mathbf{f}^{(j)}(0, t)$, where $\mathbf{f}^{(j)}(0, t)$ is the j th derivative with respect to x , evaluated at $x = 0$, we can write

$$\Delta^i \mathbf{f}(x, t) = \frac{1}{x^i} \left(\mathbf{f}(x, t) - \sum_{j=0}^{i-1} \frac{x^j}{j!} \mathbf{f}^{(j)}(0, t) \right).$$

The functional equation (3.1) can therefore be rewritten in the form

$$\left(x^k \mathbf{I}_n - t \sum_{i=0}^k x^{k-i} \mathbf{B}_i \right) \mathbf{f}(x, t) = x^k \mathbf{a} - t \sum_{j=0}^{k-1} \left(\sum_{i=j+1}^k \frac{x^{k+j-i}}{j!} \mathbf{B}_i \right) \mathbf{f}^{(j)}(0, t). \quad (3.2)$$

The key to the proof is the matrix on the left side. We will first prove the theorem under the additional assumption that this matrix has the form $x^k \mathbf{I}_n - t \mathbf{P}$ for some matrix \mathbf{P} such that $[t^0 x^0] \mathbf{P}$ is a non-singular diagonal matrix (Lemma 3). Afterwards, the general case is reduced to this situation by a perturbation argument. In the proof of Lemma 3, we will relate eigenvectors of $[t^0 x^0] \mathbf{P}$ to eigenvectors of \mathbf{P} , using the following fact.

Lemma 1. *Let $\mathbf{P} \in \mathbb{K}[x][t]^{n \times n}$ such that $\mathbf{P}_0 = [t^0] \mathbf{P} \in \mathbb{K}^{n \times n}$, and let $\mathbf{K} = x^k \mathbf{I}_n - t \mathbf{P} \in \mathbb{K}[x][t]^{n \times n}$, for some $k \in \mathbb{N}$. Let λ be an eigenvalue of \mathbf{P}_0 , let ω be a primitive k -th root of unity, and let $i \in \{0, \dots, k-1\}$. Then there is a series $y(t) = \omega^i \lambda^{1/k} t^{1/k} + \mathcal{O}(t^{2/k}) \in \mathbb{K}[[t^{1/k}]]$ such that $\det(\mathbf{K}(y(t), t)) = 0$. Furthermore, there is a vector $\mathbf{v}(t) \in \mathbb{K}[[t^{1/k}]]^n$ with algebraic coordinates such that $\mathbf{v}(t) \mathbf{K}(y(t), t) = 0$ and $\mathbf{v}(0)$ is an eigenvector of \mathbf{P}_0 for λ .*

Proof. By definition of the determinant, $\det(\mathbf{K}) = \det(x^k \mathbf{I}_n - t \mathbf{P}) = \det(x^k \mathbf{I}_n - t \mathbf{P}_0) + \mathcal{O}(t^2)$. The polynomial $\det(x^k \mathbf{I}_n - t \mathbf{P}_0) = (-1)^n t^n \det(\mathbf{P}_0 - \frac{x^k}{t} \mathbf{I}_n) \in \mathbb{K}(t)[x]$ has the root $\omega^i \lambda^{1/k} t^{1/k}$. Hence, by the Newton-Puiseux algorithm [17], there is a series $y(t) = \omega^i \lambda^{1/k} t^{1/k} + \mathcal{O}(t^{2/k}) \in \mathbb{K}[[t^{1/k}]]$ such that $\det(\mathbf{K}(y(t), t)) = 0$. It follows that the matrix $\mathbf{K}(y(t), t) \in \mathbb{K}(y(t), t)^{n \times n}$ is singular, so its left-kernel has a nonzero element $\mathbf{v} \in$

$\mathbb{K}(y(t), t)^n$. Since $y(t)$ is algebraic (because $\det(\mathbf{K})$ is a polynomial in x), also the components of \mathbf{v} are algebraic. After multiplying by a suitable power of t , we may assume that the components of $\mathbf{v}(t)$ are in $\mathbb{K}[[t^{1/k}]]$ and that $\mathbf{v}(0)$ is not the zero vector. Then $\mathbf{v}(t)\mathbf{K}(y(t), t) = 0$ implies $\mathbf{v}(0)(\lambda\mathbf{I}_n - \mathbf{P}_0) = 0$, which completes the proof. \square

Secondly, we will need to ensure that a certain matrix is nonsingular.

Lemma 2. *Let $\lambda_0, \dots, \lambda_{n-1} \in \mathbb{K} \setminus \{0\}$ and let ω be a primitive k -th root of unity. For $u, v = 0, \dots, nk - 1$ define $c_{u,v} = (\omega^{u \bmod k} \lambda_{\lfloor u/k \rfloor})^{\lfloor v/n \rfloor} \delta_{\lfloor u/k \rfloor, \lfloor v/k \rfloor}$. Then*

$$\det((c_{u,v})_{u,v=0}^{nk-1}) = \pm \left(\prod_{0 \leq i < j < k} (\omega^j - \omega^i) \right)^n \prod_{\ell=0}^{n-1} \lambda_{\ell}^{\binom{k}{2}}.$$

In particular, this determinant is not zero.

Proof. The expression on the right is non-zero because $\lambda_{\ell} \neq 0$ for all ℓ and ω is a primitive root of unity. It remains to prove the claimed identity.

We permute the columns of the matrix such that the entry at position (u, v) is

$$(\omega^{u \bmod k} \lambda_{\lfloor u/k \rfloor})^{v \bmod k} \delta_{\lfloor u/k \rfloor, \lfloor v/k \rfloor}.$$

The resulting matrix is block diagonal with n blocks of size $k \times k$. The ℓ -th block is the Vandermonde matrix $((\omega^{ij} \lambda_{\ell}^j))_{i,j=0}^{k-1}$, whose determinant is $\lambda_{\ell}^{\binom{k}{2}} \prod_{i < j} (\omega^j - \omega^i)$. Since the determinant of a block diagonal matrix is the product of the determinants of its blocks, we arrive at the desired conclusion. \square

The idea for the proof of Lemma 3 is to replace x by various algebraic series $y(t)$ in such a way that the terms $\mathbf{f}(x, t)$ in (3.2) disappear and a linear system for the components of $\mathbf{f}^{(j)}(0, t)$ arises, and then to show that this system has a unique solution.

Lemma 3. *Let $\lambda_0, \dots, \lambda_{n-1} \in \mathbb{K} \setminus \{0\}$ be pairwise distinct, and let $\mathbf{E} = \text{diag}(\lambda_0, \dots, \lambda_{n-1}) \in \mathbb{K}^{n \times n}$. Let $\mathbf{a} \in \mathbb{K}[x][t]^n$, and let $\mathbf{P}, \mathbf{Q}_0, \dots, \mathbf{Q}_{k-1} \in \mathbb{K}[x][t]^{n \times n}$. Suppose that $\mathbf{P} = \mathbf{E} + \mathbf{O}(t)$ and $\mathbf{Q}_j = x^j \mathbf{E} + \mathbf{O}(t)$ for $j = 0, \dots, k-1$. If $\mathbf{f} \in \mathbb{K}[x][[t]]^n$ and $\mathbf{g}_0, \dots, \mathbf{g}_{k-1} \in \mathbb{K}[[t]]^n$ are such that*

$$(x^k \mathbf{I}_n - t \mathbf{P}) \mathbf{f} = x^k \mathbf{a} - t \sum_{j=0}^{k-1} \mathbf{Q}_j \mathbf{g}_j, \quad (3.3)$$

then the components of $\mathbf{g}_0, \dots, \mathbf{g}_{k-1}$ and \mathbf{f} are algebraic over $\mathbb{K}[x][t]$.

Proof. By Lemma 1, the polynomial $\det(x^k \mathbf{I}_n - t \mathbf{P})$ has nk series roots $y_{ij}(t)$ of the form $y_{ij}(t) = \omega^i \lambda_j^{1/k} t^{1/k} + \mathbf{O}(t^{2/k})$. For each such root, the matrix $y_{ij}(t)^k \mathbf{I}_n - t \mathbf{P}(y_{ij}(t), t)$ is

singular, and, again by Lemma 1, the left-kernel of $y_{ij}(t)^k \mathbf{I}_n - t\mathbf{P}(y_{ij}(t), t)$ contains a vector $\mathbf{v}_{ij}(t) = \lambda_j^{-1} \mathbf{e}_j + \mathcal{O}(t^{1/k})$, with algebraic coefficients. Here \mathbf{e}_j is the j th unit vector.

For $i = 0, \dots, k-1$ and $j = 0, \dots, n-1$ we replace x by $y_{ij}(t)$ in equation (3.3) and multiply with $\mathbf{v}_{ij}(t)$ from the left. This gives an inhomogeneous linear system with nk equations for the nk unknown components of $\mathbf{g}_0(t), \dots, \mathbf{g}_{k-1}(t) \in \mathbb{K}[[t]]^n$, whose coefficient matrix is

$$\begin{pmatrix} \mathbf{v}_{0,0}(t)\mathbf{Q}_0(y_{0,0}(t), t) & \cdots & \mathbf{v}_{0,0}(t)\mathbf{Q}_{k-1}(y_{0,0}(t), t) \\ \mathbf{v}_{1,0}(t)\mathbf{Q}_0(y_{1,0}(t), t) & \cdots & \mathbf{v}_{1,0}(t)\mathbf{Q}_{k-1}(y_{1,0}(t), t) \\ \vdots & \ddots & \vdots \\ \mathbf{v}_{k-1,n-1}(t)\mathbf{Q}_0(y_{k-1,n-1}(t), t) & \cdots & \mathbf{v}_{k-1,n-1}(t)\mathbf{Q}_{k-1}(y_{k-1,n-1}(t), t) \end{pmatrix}.$$

We are done if we can show that this matrix is invertible, because this implies that the inhomogeneous linear system for the components of $\mathbf{g}_0, \dots, \mathbf{g}_{k-1}$ has a unique solution. The components of its solution vector must be algebraic, because all the series appearing in the linear system are algebraic. From equation (3.3), we finally see that the algebraicity of the components of $\mathbf{g}_0, \dots, \mathbf{g}_{k-1}$ implies the algebraicity of the components of \mathbf{f} .

To see that the matrix above is invertible, we use the assumption that $\mathbf{Q}_\ell(y_{ij}(t), t) = \mathbf{E}y_{ij}(t)^\ell + \mathcal{O}(t) = (\omega^i \lambda_j^{1/k})^\ell \mathbf{E}t^{\ell/k} + \mathcal{O}(t^{(\ell+1)/k}) \in \mathbb{K}[[t^{1/k}]]^{n \times n}$. Together with $\mathbf{e}_j \mathbf{E} = \lambda_j \mathbf{e}_j$, where \mathbf{e}_j is again the j th unit vector, it follows that

$$\mathbf{v}_{ij}(t)\mathbf{Q}_\ell(y_{ij}(t), t) = (\omega^i \lambda_j^{1/k})^\ell \mathbf{e}_j t^{\ell/k} + \mathcal{O}(t^{(\ell+1)/k}) \in \mathbb{K}[[t^{1/k}]]^n,$$

for $i = 0, \dots, k-1$, $j = 0, \dots, n-1$ and $\ell = 0, \dots, k-1$. Therefore, for $u = 0, \dots, nk-1$ and $v = 0, \dots, nk-1$, the entry of the matrix at position (u, v) is

$$c_{u,v} := (\omega^{u \bmod k} \lambda_{\lfloor u/k \rfloor}^{1/k})^{\lfloor v/n \rfloor} t^{\lfloor v/n \rfloor / k} \delta_{\lfloor u/k \rfloor, v \bmod n} + \mathcal{O}(t^{(\lfloor v/n \rfloor + 1)/k}).$$

By Lemma 2, we have $\det((c_{u,v})_{u,v=0}^{nk-1}) \neq 0$. This completes the proof. \square

Proof of Thm. 2. We have already argued that existence and uniqueness of a solution in $\mathbb{K}[x][[t]]^n$ are evident. To show that its components are algebraic, we bring equation (3.1) into a form where Lemma 3 applies. Let ϵ be a new variable, let $\lambda_1, \dots, \lambda_n \in \mathbb{K} \setminus \{0\}$ be pairwise distinct and set $\mathbf{E} = \epsilon \operatorname{diag}(\lambda_1, \dots, \lambda_n)$. Set $\tilde{\mathbf{a}}(x, t) := \mathbf{a}(x, t^2)$, $\tilde{\mathbf{B}}_i(x, t) := t\mathbf{B}_i(x, t^2)$ ($i = 0, \dots, k-1$), and $\tilde{\mathbf{B}}_k(x, t) := \mathbf{E} + t\mathbf{B}_k(x, t^2)$, and consider the system

$$\tilde{\mathbf{f}} = \tilde{\mathbf{a}} + t \sum_{i=0}^k \tilde{\mathbf{B}}_i \Delta^i \tilde{\mathbf{f}}.$$

This system has a unique solution $\tilde{\mathbf{f}} \in \mathbb{K}[x, \epsilon][[t]]^n$, which is related to the solution \mathbf{f} of the original equation (3.1) via $\mathbf{f}(x, t^2) = [\epsilon^0] \tilde{\mathbf{f}}(x, t)$. We are done if we can show that the components of $\tilde{\mathbf{f}}$ are algebraic, because then so are the components of \mathbf{f} .

Indeed, translating the delta terms to partial derivatives, as earlier, gives

$$\left(x^k \mathbf{I}_n - t \sum_{i=0}^k x^{k-i} \tilde{\mathbf{B}}_i\right) \tilde{\mathbf{f}}(x, t) = x^k \tilde{\mathbf{a}} - t \sum_{j=0}^{k-1} \left(\sum_{i=j+1}^k \frac{x^{k+j-i}}{j!} \tilde{\mathbf{B}}_i\right) \tilde{\mathbf{f}}^{(j)}(0, t).$$

For the matrix $\mathbf{P} = \sum_{i=0}^k x^{k-i} \tilde{\mathbf{B}}_i \in \mathbb{K}(\epsilon)[x, t]^{n \times n}$ we have $\mathbf{P} = \mathbf{E} + \mathcal{O}(t)$, and for the matrices $\mathbf{Q}_j = k! \sum_{i=j+1}^k \frac{x^{k+j-i}}{j!} \tilde{\mathbf{B}}_i \in \mathbb{K}(\epsilon)[x, t]^{n \times n}$ we have $\mathbf{Q}_j = x^j \mathbf{E} + \mathcal{O}(t)$ for $j = 0, \dots, k-1$. Therefore, Lemma 3 applies to the perturbed equation above and yields the desired algebraicity result. (The lemma is applied with \mathbb{K} replaced by some algebraic closure of $\mathbb{K}(\epsilon)$ and with $\tilde{\mathbf{f}}^{(j)}(0, t)/k!$ in the role of \mathbf{g}_j .) \square

4 Models with more than one restriction

We have seen that inhomogeneous models in a half-space $\mathbb{Z}_{\geq 0} \times \mathbb{Z}^{d-1}$ always have an algebraic generating function. More generally, consider walks restricted to $\mathbb{Z}_{\geq 0}^{d_1} \times \mathbb{Z}^{d_2}$. In this case, the question arises whether some of the d_1 constraints are implied by the others, which has led Bostan et al. [10] to introduce the *dimension* of a model. Here we generalize this notion to inhomogeneous walks.

First consider unrestricted models in \mathbb{Z}^d . Fix an inhomogeneity as in Section 2. Let \mathbf{S} be the union of some disjoint copies of the sets \mathbf{S}_{pq} , so that a walk in \mathbb{Z}^d of length n can be viewed as a word w over the alphabet \mathbf{S} . To any such walk w , we associate the vector $(a_{\mathbf{u}})_{\mathbf{u} \in \mathbf{S}}$ where $a_{\mathbf{u}} \in \mathbb{N}$ is the number of occurrences of \mathbf{u} in w . While for unrestricted homogeneous models, every vector of natural numbers is associated with some walk, this is not true for inhomogeneous models. For example, for space-inhomogeneous walks in \mathbb{Z}^2 with $\mathbf{S}_0 = \{\nearrow\}$ and $\mathbf{S}_1 = \{\swarrow\}$, the vector $(1, 1)$ is not associated with a walk. The next lemma is a characterization of vectors associated with walks.

Lemma 4. *Consider a finite automaton and the inhomogeneity defined by it as in Section 2. Let \mathbf{S} be a disjoint union of the sets \mathbf{S}_{pq} , and $(a_{\mathbf{u}})_{\mathbf{u} \in \mathbf{S}}$ be a vector of nonnegative integers. Let G be the multigraph obtained from the automaton by replacing an edge labelled with \mathbf{u} by $a_{\mathbf{u}}$ many (unlabelled) edges. Then $(a_{\mathbf{u}})_{\mathbf{u} \in \mathbf{S}}$ is associated with a walk if and only if G has an Eulerian path starting at the initial state q_0 and ending at some final state.*

Proof. Since \mathbf{S} is the union of disjoint copies of the step sets \mathbf{S}_{pq} , every step $\mathbf{u} \in \mathbf{S}$ belongs to exactly one such set. Therefore, any walk w in the model which has $(a_{\mathbf{u}})_{\mathbf{u} \in \mathbf{S}}$ as associated vector can be translated into a path in G starting at q_0 , ending at some final state, and using $\sum_{\mathbf{u} \in \mathbf{S}_{pq}} a_{\mathbf{u}}$ times an edge from p to q , for all vertices p and q . This is an Eulerian path. Conversely, let (q_0, q_1, \dots, q_n) be an Eulerian path in G with q_n being a final state. Then for any two p, q states there are $\sum_{\mathbf{u} \in \mathbf{S}_{pq}} a_{\mathbf{u}}$ many indices $i \in \{1, \dots, n\}$ such that $(q_{i-1}, q_i) = (p, q)$. For every $i \in \{1, \dots, n\}$, assign an arbitrary step $\mathbf{u} \in \mathbf{S}_{q_{i-1}q_i}$

in such that each step \mathbf{u} is chosen exactly $a_{\mathbf{u}}$ times, and let w be the walk composed of the selected steps. By definition of the \mathbf{S}_{pq} 's, it belongs to the model. \square

The condition for a graph to have an Eulerian path can be encoded as a system of linear constraints on the in-degree and out-degrees of the vertices of the graph [6]. In our case these are linear equations for the variables $a_{\mathbf{u}}$. It can also be encoded into linear equations that the Eulerian path should start or end at prescribed vertices.

A walk w ends in $\mathbb{Z}_{\geq 0}^{d_1} \times \mathbb{Z}^{d_2}$ iff for its associated vector $(a_{\mathbf{u}})_{\mathbf{u} \in \mathbf{S}}$ we have

$$\sum_{\mathbf{u}=(u_1, \dots, u_{p+q}) \in \mathbf{S}} a_{\mathbf{u}} u_i \geq 0 \quad \text{for all } i = 1, \dots, d_1, \quad (4.1)$$

and it stays entirely in $\mathbb{Z}_{\geq 0}^{d_1} \times \mathbb{Z}^{d_2}$ iff these inequalities hold for all prefixes of w . Extending Def. 2 of [10], we define the *dimension* of an inhomogeneous model for $\mathbb{Z}_{\geq 0}^{d_1} \times \mathbb{Z}^{d_2}$ as the smallest $\delta \in \mathbb{N}$ such that there are δ inequalities in (4.1) which imply all the others. In view of Lemma 4, the dimension of a model can be found by linear programming.

5 Inhomogeneous Lattice Walks in the Quarter Plane

We have no satisfactory theory for inhomogeneous models for the quarter plane. However, for all time-inhomogeneous and all space-inhomogeneous models with small steps as in the introduction, we have produced an experimental classification which is available at <http://www.algebra.uni-linz.ac.at/people/mkauers/inhomogeneous/>. Up to symmetry, there are 32993 pairs $\mathbf{S}_0, \mathbf{S}_1 \subseteq \mathbb{N}^2$. Removing trivial cases (whose counting sequence is ultimately constant), zero- and one-dimensional cases (whose generating function is algebraic by Thm. 1), and homogeneous cases (whose nature is known) leaves us with 23906 space-inhomogeneous and 25370 time-inhomogeneous cases.

For each of these, we computed the first 10000 terms (modulo the prime 45007) of the generating function $F(1, 1, t)$ and tried to guess a differential equation. When an equation was found, we also searched for an algebraic equation. These computations took altogether about 30 years of computing time. As a result, 3784 space-inhomogeneous models and 2603 time-inhomogeneous models seem to be D-finite, including 2474 and 1535 seemingly algebraic cases, respectively. For space-inhomogeneous models, the largest differential equations we found have order 24 and degree 1183, such an equation appears for $\mathbf{S}_0 = \nearrow$ and $\mathbf{S}_1 = \nwarrow$. For time-inhomogeneous models, the largest differential equation we found has order 28 and degree 1256 and appears for $\mathbf{S}_0 = \leftrightarrow$, $\mathbf{S}_1 = \nleftrightarrow$. Very likely further D-finite models could be discovered with more terms.

The techniques of [12] for proving D-finiteness seem to apply only to a very limited number of cases. We conclude with two examples where they work and invite our readers to find proofs for further conjecturally D-finite cases.

Example 4. Consider the time-inhomogeneous model in $\mathbb{Z}_{\geq 0}^2$ with $\mathbf{S}_0 = \begin{smallmatrix} \swarrow \\ \downarrow \\ \searrow \end{smallmatrix}$ and $\mathbf{S}_1 = \begin{smallmatrix} \swarrow & \searrow \\ \downarrow & \downarrow \\ \swarrow & \searrow \end{smallmatrix}$. Let $S_0, S_1 \in \mathbb{Q}[x, x^{-1}, y, y^{-1}]$ be the corresponding Laurent polynomials, and let $F_0(x, y), F_1(x, y) \in \mathbb{Q}[x, y][[t]]$ be the power series counting the number of walks of even and odd lengths, respectively. Then $F(x, y) = F_0(x, y) + F_1(x, y)$ is the generating function of the model. The functional equations for $F_0(x, y)$ and $F_1(x, y)$ are

$$\begin{aligned} F_0(x, y) &= 1 + tS_1F_1(x, y) - t([y^{<0}]S_1)F_1(x, 0) - t([x^{<0}]S_1)F_1(0, y) + t([x^{<0}y^{<0}]S_0)F_1(0, 0) \\ F_1(x, y) &= tS_0F_0(x, y) - t([y^{<0}]S_0)F_0(x, 0). \end{aligned}$$

We consider the groups G_0 and G_1 generated by the rational maps $\Phi_0: (x, y) \mapsto (\frac{1}{x}, y)$ and $\Psi_0: (x, y) \mapsto (x, \frac{1}{y(x + \frac{1}{x})})$, and $\Phi_1: (x, y) \mapsto (\frac{1}{x}, y)$ and $\Psi_1: (x, y) \mapsto (x, \frac{1}{y(x + \frac{1}{x})})$, respectively. These groups act on pairs of rational functions. By construction their generators alter only one of two of its components, and leave the corresponding step polynomials invariant, see [12] for details. We multiply the two equations above by xy and take the so-called orbit sum of the first equation with respect to G_1 , and of the second one with respect to G_0 . This eliminates all terms $F_q(x, 0)$, $F_q(0, y)$ and $F_q(0, 0)$ with $q \in \{0, 1\}$ and leaves us with

$$\begin{aligned} \sum_{g \in G_1} \text{sgn}(g)g(xyF_0(x, y)) &= \sum_{g \in G_1} \text{sgn}(g)g(xy) + tS_1 \sum_{g \in G_1} \text{sgn}(g)g(xyF_1(x, y)) \\ \sum_{g \in G_0} \text{sgn}(g)g(xyF_1(x, y)) &= tS_0 \sum_{g \in G_0} \text{sgn}(g)g(xyF_0(x, y)), \end{aligned}$$

where $\text{sgn}(g) = (-1)^\ell$ when g is a product of ℓ generators. It is easy to check that replacing y by $\frac{1}{y(x + \frac{1}{x})}$ in the second equation gives

$$\sum_{g \in G_1} \text{sgn}(g)g(xyF_1(x, y)) = tS_0(x, \frac{1}{y(x + \frac{1}{x})}) \sum_{g \in G_1} \text{sgn}(g)g(xyF_0(x, y)).$$

From this equation and the first of the two previous two equations we get

$$\sum_{g \in G_1} \text{sgn}(g)g(xyF_0(x, y)) = \frac{1}{1 - t^2S_0(x, \frac{1}{y(x + \frac{1}{x})})S_1} \sum_{g \in G_1} \text{sgn}(g)g(xy)$$

Extracting the positive part gives

$$F_0(x, y) = \frac{1}{xy} [x^{>0}y^{>0}] \frac{1}{1 - t^2S_0(x, \frac{1}{y(x + \frac{1}{x})})S_1} \sum_{g \in G_1} \text{sgn}(g)g(xy).$$

This expression implies D -finiteness of $F_0(x, y)$, and back-substituting into the earlier equations and using D -finite closure properties gives the D -finiteness of $F_1(x, y)$ and of $F(x, y)$.

Example 5. Consider the space-inhomogeneous model in $\mathbb{Z}_{\geq 0}^2$ with $\mathbf{S}_0 = \leftrightarrow$ and $\mathbf{S}_1 = \bowtie$ studied by D'Arco et al. [15]. Define \mathbf{S}_{pq} for $p, q \in \{0, 1\}$ as in Section 2, and let $S_{pq} \in \mathbb{Q}[x, x^{-1}, y, y^{-1}]$ be the corresponding Laurent polynomials. Write $F_q(x, y)$ for $q \in \{0, 1\}$ for the power series that count the number of walks ending at points (i, j) with $i + j = q \pmod 2$. The functional equations for $F_0(x, y)$ and $F_1(x, y)$ are

$$\begin{aligned} F_0(x, y) &= 1 + tS_{10}F_1(x, y) - t([y^{<0}]S_{10})F_1(x, 0) - t([x^{<0}]S_{10})F_1(0, y) \\ F_1(x, y) &= tS_{01}F_0(x, y) - t([y^{<0}]S_{01})F_0(x, 0) - t([x^{<0}]S_{01})F_0(0, y) \\ &\quad + tS_{11}F_1(x, y) - t([y^{<0}]S_{11})F_1(x, 0) - t([x^{<0}]S_{11})F_1(0, y). \end{aligned}$$

Consider the group G generated by the rational maps $\Phi: (x, y) \mapsto (x, \frac{1}{y})$ and $\Psi: (x, y) \mapsto (\frac{1}{x}, y)$. Like in the previous example, multiply both equations by xy and take the orbit sum for G . This eliminates all terms $F_q(x, 0)$ and $F_q(0, y)$ and leaves us with

$$\begin{aligned} \sum_{g \in G} \text{sgn}(g)g(xyF_0(x, y)) &= \sum_{g \in G} \text{sgn}(g)g(xy) + tS_{10} \sum_{g \in G} \text{sgn}(g)g(xyF_1(x, y)) \\ \sum_{g \in G} \text{sgn}(g)g(xyF_1(x, y)) &= tS_{01} \sum_{g \in G} \text{sgn}(g)g(xyF_0(x, y)) + tS_{11} \sum_{g \in G} \text{sgn}(g)g(xyF_1(x, y)). \end{aligned}$$

From those equations we deduce

$$\sum_{g \in G} \text{sgn}(g)g(xyF_1(x, y)) = \frac{tS_{01}}{1 - tS_{11} - t^2S_{01}S_{10}} \sum_{g \in G} \text{sgn}(g)g(xy),$$

and extracting the positive part gives

$$F_1(x, y) = \frac{1}{xy} [x^{>0}y^{>0}] \frac{tS_0}{1 - tS_{11} - t^2S_{01}S_{10}} \sum_{g \in G} \text{sgn}(g)g(xy).$$

This expression implies D -finiteness of $F_1(x, y)$, and back-substituting into the earlier equations and using D -finite closure properties gives the D -finiteness of $F_0(x, y)$ and of the full generating function $F(x, y) = F_0(x, y) + F_1(x, y)$.

We were able to guess a differential equation of order 3 with polynomial coefficients of degree 9 for the generating function $F(1, 1, t)$ in Example 5. For the model in Example 4 we could not find any.

Acknowledgement. We thank Marni Mishna for making us aware of [13], and the referees for pointing us to [19].

References

- [1] Axel Bacher, Manuel Kauers, and Rika Yatchak. “Continued Classification of 3D Lattice Walks in the Positive Octant”. In: *Proceedings of FPSAC'16*. 2016, pp. 95–106.

- [2] Cyril Banderier and Philippe Flajolet. “Basic Analytic Combinatorics of Directed Lattice Paths”. In: *Theoretical Computer Science* 281.1–2 (2002), pp. 37–80.
- [3] N. R. Beaton, A. L. Owczarek, and A. Rechnitzer. *Exact solution of some quarter plane walks with interacting boundaries*. Tech. rep. 1807.08853. ArXiv, 2018.
- [4] Olivier Bernardi, Mireille Bousquet-Mélou, and Kilian Raschel. *Counting quadrant walks via Tutte’s invariant method*. Tech. rep. 1708.08215. ArXiv, 2017.
- [5] Benjamin Bogosel et al. *3D positive lattice walks and spherical triangles*. Tech. rep. 1807.08610. ArXiv, 2018.
- [6] Béla Bollobás. *Modern Graph Theory*. Springer, 1998.
- [7] Nicolas Bonichon et al. “On the number of planar Eulerian orientations”. In: *European Journal of Combinatorics* 65 (2016), pp. 59–91.
- [8] Alin Bostan. *Calcul Formel pour la Combinatoire des Marches*. Habilitation a Diriger des Recherches, Université Paris 13. 2017.
- [9] Alin Bostan, Mireille Bousquet-Mélou, and Stephen Melczer. *Counting walks with large steps in an orthant*. Tech. rep. 1806.00968. ArXiv, 2018.
- [10] Alin Bostan et al. “On 3-dimensional lattice walks confined to the positive octant”. In: *Annals of Combinatorics* 20.4 (2016), pp. 661–704.
- [11] Mireille Bousquet-Mélou and Arnaud Jehanne. “Polynomial equations with one catalytic variable, algebraic series and map enumeration”. In: *Journal of Combinatorial Theory, Series B* 96.5 (2006), pp. 623–672.
- [12] Mireille Bousquet-Mélou and Marni Mishna. “Walks with small steps in the quarter plane”. In: *Contemporary Mathematics* 520 (2010), pp. 1–40.
- [13] Mireille Bousquet-Mélou and Guoce Xin. “On partitions avoiding 3-crossings”. In: *Séminaire Lotharingien de Combinatoire* 54 (2006).
- [14] Julien Courtiel et al. “Weighted Lattice Walks and Universality Classes”. In: *Journal of Combinatorial Theory, Series A* 152 (2017), pp. 255–302.
- [15] Philippe D’Arco, Valentiana Lacivita, and Sami Mustapha. “Combinatorics meets potential theory”. In: *Electronic Journal of Combinatorics* 23.2 (2018), P2.28.
- [16] Thomas Dreyfus et al. “On the Nature of the Generating Series of Walks in the Quarter Plane”. In: *Inventiones mathematicae* 213.1 (2018), pp. 205–236.
- [17] Manuel Kauers and Peter Paule. *The Concrete Tetrahedron*. Springer, 2011.
- [18] Manuel Kauers and Rika Yatchak. “Walks in the Quarter Plane with Multiple Steps”. In: *Proceedings of FPSAC’15*. DMTCS. 2015, pp. 35–36.
- [19] Dorin Popescu. “General Néron desingularization and approximation”. In: *Nagoya Mathematical Journal* 104 (1986), pp. 85–115.