

Application of Unspecified Sequences in Symbolic Summation

Manuel Kauers^{*}
RISC-Linz
Johannes Kepler Universität
A-4040 Linz, Austria
mkauers@risc.uni-linz.ac.at

Carsten Schneider^{*}
RISC-Linz
Johannes Kepler Universität
A-4040 Linz, Austria
cschneid@risc.uni-linz.ac.at

ABSTRACT

We consider symbolic sums which contain subexpressions representing unspecified sequences. Existing symbolic summation technology is extended to sums of this kind. We show how this can be applied in the systematic search for general summation identities. Both, results about the non-existence of identities of a certain form, and examples of general families of identities which we have discovered automatically are included in the paper.

Categories and Subject Descriptors

I.1.2 [Computing Methodologies]: Symbolic and Algebraic Manipulation—*Algorithms*; G.2.1 [Discrete Mathematics]: Combinatorics—*Recurrences and difference equations*

General Terms

Algorithms

Keywords

Symbolic Summation, Difference Fields

1. INTRODUCTION

The focus of this paper is on summation identities involving expressions like X_k that stand for *unspecified sequences* (or *generic sequences*). Such summation identities remain true for any interpretation of X_k as a particular sequence, i.e., we may put $X_k = f_k$ for any sequence $(f_k)_{k \geq 0}$. A simple example for such an identity is

$$\sum_{k=0}^n (X_{k+1} - 1) \prod_{i=0}^k X_i = \prod_{k=0}^{n+1} X_k - X_0. \quad (1)$$

^{*}Both authors were supported by the Austrian science foundation FWF, grants P16613-N12 and F1305.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.

ISSAC'06, July 9–12, 2006, Genova, Italy.

Copyright 2006 ACM 1-59593-276-3/06/0004 ...\$5.00.

If particular rational functions are substituted for X_k in this identity, we obtain precisely the indefinite hypergeometric summation identities that are found by Gosper's algorithm [9, 16]. (Subtle exceptions may arise owing to the singularities of a particular rational function, see [4, 2]. These are ignored throughout the present paper.) We may say that Gosper's algorithm precisely solves the task of writing a given hypergeometric term $\prod_{i=0}^k X_k$ in the form $(Y_{k+1} - 1) \prod_{i=0}^k Y_i$ for a specific rational function Y_k , or it proves that this is impossible.

In the present paper, we mainly study definite summation identities involving unspecified sequences. A sum is called definite if the summand depends not only on the summation index k but also on the summation bound n . A simple example is

$$\sum_{k=0}^n X_{n+k} = X_0 + \sum_{k=0}^{n-1} (X_{2k+1} + X_{2k+2} - X_k)$$

Again, the identity holds for every sequence in place of the symbol X_k . The interest in identities of this type was not so much raised by examples arising from practice. The consideration of summation identities involving unspecified sequences X_k is rather motivated because the presence of X_k in summation identities reveals some structural information about summation in general and summation algorithms in particular. The use of unspecified sequences makes it possible to search for (families of) “nice” summation identities in a more systematic way. For instance, it is possible to automatically derive criteria for finding particular sequences $(f_k)_{k \geq 0}$, which, in place of X_k , lead to a further simplification of the sum.

An earlier paper of ours [12] contains an algorithm for indefinite summation with unspecified sequences. In Section 2 we discuss how this algorithm can be extended to definite summation via the creative telescoping method [26]. This algorithm was implemented and incorporated into the second author's summation package Sigma [22]. With this implementation, we searched for general definite summation identities involving unspecified sequences. It turned out that only very few general identities exist. An explanation for this phenomenon is provided in Section 3, where we show that a certain class of definite sums does not admit creative telescoping recurrences at all.

This negative result suggests that the search for nontrivial general summation identities has to be focused on more complicated expressions, such as nested sum expressions. Even

in this enlarged domain, general definite sums which admit simplification are rare. Some new identities, which we have found by experimenting, are presented in Section 4.

2. SUMMATION IN DIFFERENCE FIELDS

2.1 PLDEs and Summation

A fundamental role in summation algorithms is played by *parameterized linear difference equations* (PLDEs): Given certain sequences a_0, \dots, a_m and $f_0, \dots, f_r: \mathbb{N} \rightarrow \mathbb{K}$, where \mathbb{K} is a field, the goal is to find a sequence g and constants c_0, \dots, c_r such that

$$a_0g + a_1Eg + \dots + a_mE^mg = c_0f_0 + \dots + c_rf_r, \quad (2)$$

where E denotes the shift operator $(Ef)_k := f_{k+1}$.

Indefinite summation provides the most special situation. Here, we aim at eliminating the summation quantifier from $\sum_{k=0}^n f_k$ with f_k independent of n . If g is a solution of (2) with $m = 1$, $r = 0$, $a_0 = -1$, $a_1 = 1$, $f_0 = f$ (telescoping equation), then we have $\sum_{k=0}^n f_k = g_{n+1} - g_0$ (with $c_0 = 1$).

Less straightforward are definite sums. These are sums of the form $S_n = \sum_{k=0}^n f_{n,k}$, where $f_{n,k}$ may depend on both, the summation index k and the bound n . For such sums, telescoping normally fails. Therefore, we try a different strategy. We proceed in two steps: First, we compute a recurrence equation for the sum by means of *creative telescoping* [26], and in a second step we solve that recurrence. In creative telescoping, we consider k as the independent variable and n as constant, and we solve

$$Eg - g = c_0f_0 + c_1f_1 + \dots + c_rf_r \quad (3)$$

with $f_i = f_{n+i,k}$ (another special case of (2)). If no solution exists, we increase r and try again. Once a solution (c_0, \dots, c_r, g) is found, we can derive a recurrence for the original sum S by summing the equation over k . This recurrence is of the form

$$a_0S + a_1ES + \dots + a_rE^rS = f$$

(another special case of (2)) for certain a_0, \dots, a_r, f which originate from c_0, \dots, c_r , and g in (3). Now n is considered as independent variable, and k is no longer present in a_i or f . Solving this equation for S delivers an explicit representation for the definite sum.

This general summation strategy is explained in detail for hypergeometric summation in [19]. Implementations of these algorithms are available since long [18, 1]. The same technique is applicable for more general expressions.

Example 1. Consider the definite sum $S_n = \sum_{k=0}^n f_{n,k}$ with $f_{n,k} = kX_{n+k}$, where X_{n+k} denotes a generic sequence. Creative telescoping is applied for $r = 0, 1, 2, 3, \dots$ in turn, until a solution is found. Choosing, for instance, $r = 2$, we have to find constants c_0, c_1, c_2 and some $g_{n,k}$ such that

$$\begin{aligned} g_{n,k+1} - g_{n,k} &= c_0f_{n,k} + c_1f_{n+1,k} + c_2f_{n+2,k} \\ &= c_0kX_{n+k} + c_1kX_{n+k+1} + c_2kX_{n+k+2}. \end{aligned}$$

It is easily checked that a solution is given by $(c_0, c_1, c_2, g) = (1, -2, 1, (k-1)X_{n+k+1} - kX_{n+k})$. Summing the equation over k from 0 to n and compensating for the missing terms gives the recurrence

$$S_n - 2S_{n+1} + S_{n+2} = f_n$$

with

$$\begin{aligned} f_n &= -(-X_{n+1} + (n+1)X_{2n+1} \\ &\quad + (n+2)X_{2n+2} - (n+1)X_{2n+3} - (n+2)X_{2n+4}). \end{aligned}$$

Next, we have to solve this recurrence. The homogeneous equation obviously has the solutions 1 and n , and it turns out that the inhomogeneous equation has $\sum_{k=1}^n \sum_{i=1}^k f_{i-2}$ as a particular solution. Comparison of two initial values reveals the representation

$$S_n = \sum_{k=1}^n \sum_{i=1}^k f_{i-2}. \quad (4)$$

This can be simplified further, as we will see below (Example 3).

This summation process can be carried out computationally in the general setting of *difference fields* [22]. A difference field is a pair (\mathbb{F}, σ) where $\sigma: \mathbb{F} \rightarrow \mathbb{F}$ is a field automorphism. The elements of a difference field are understood as formalizations of sequences, and the automorphism σ should act on the field elements like the shift operator $(Ef)_k = f_{k+1}$ acts on the corresponding sequences. The elements $c \in \mathbb{F}$ with $\sigma(c) = c$ form a subfield \mathbb{K} of \mathbb{F} , called the *field of constants*. In the language of difference fields, the problem of solving parameterized linear difference equations reads as follows:

GIVEN: A difference field (\mathbb{F}, σ) with constant field \mathbb{K} and elements $a_0, \dots, a_m \in \mathbb{F}$ and $f_0, \dots, f_r \in \mathbb{F}$.

FIND: All tuples $(c_0, \dots, c_r, g) \in \mathbb{K}^{r+1} \times \mathbb{F}$ such that

$$a_0g + a_1\sigma(g) + \dots + a_m\sigma^m(g) = c_0f_0 + \dots + c_rf_r. \quad (5)$$

It is easy to check that all the solutions (c_0, \dots, c_r, g) of a parameterized linear difference equation form a vector space over \mathbb{K} , and we want to compute a basis of this vector space. Of course, it might be that there do not exist solutions in the given field \mathbb{F} . In this case, it is of interest to construct a bigger field where there exists a “nice” solution. This will be used in Section 4 for deriving criteria on the X_k that make a given sum summable in closed form.

Algorithms for solving PLDEs are available for several types of difference fields (\mathbb{F}, σ) . For the simplest case of a constant field, i.e., $\mathbb{F} = \mathbb{K}$, the solution of a PLDE is immediate by linear algebra. In the remainder of this section, we will outline solution algorithms for free difference fields and $\Pi\Sigma$ -extensions. Difference fields constructed from free difference fields by $\Pi\Sigma$ -extensions are the appropriate fields for the summation problems we want to consider later in this paper.

2.2 Solving PLDEs in free difference fields

All difference fields used in this paper are constructed as (iterated) difference field extensions over some constant field \mathbb{K} . A difference field extension of some difference field (\mathbb{G}, σ) is a difference field (\mathbb{E}, σ') where \mathbb{G} is a subfield of \mathbb{E} and $\sigma'|_{\mathbb{G}} = \sigma$. In this paper, all difference field extensions will be obtained by adjoining one or more transcendental elements t, t', t'', \dots to the ground field \mathbb{G} and extending the definition of the shift σ to this extended field $\mathbb{G}(t, t', t'', \dots)$.

Following Cohn [6], we define the *free* difference field extension of a difference field (\mathbb{F}, σ) (by a *difference variable* x) to be the field

$$\mathbb{F}\langle x \rangle := \mathbb{F}(\dots, x_{-2}, x_{-1}, x_0, x_1, x_2, \dots).$$

The shift σ is extended from \mathbb{F} to $\mathbb{F}\langle x \rangle$ by the definition $\sigma(x_i) := x_{i+1}$ ($i \in \mathbb{Z}$). We understand here that the set $\{x_i : i \in \mathbb{Z}\}$ is algebraically independent over \mathbb{F} . A difference variable x is an appropriate algebraic representation of an unspecified sequence X_k [11]. By capital letters (X_k , Y_{k+3} , etc.) we denote subexpressions which appear in a symbolic sum and which denote an unspecified sequence. The corresponding lowercase letters (x_0 , y_3 , etc.) will be used in the corresponding difference field representation.

Difference fields which are constructed by iterated free extensions of the constant field \mathbb{K} , e.g., $\mathbb{F} = \mathbb{K}\langle x \rangle \langle y \rangle$, are called *free difference fields*. Free difference fields are very easy to deal with computationally, because for each $f \in \mathbb{F}\langle x \rangle \setminus \mathbb{F}$, the forward and backward shift $\sigma(f)$ and $\sigma^{-1}(f)$ both must contain some variable x_i which does not appear in f itself. This makes it possible to devise a solving algorithm based on simple cancellation considerations. We have already described such an algorithm in an earlier paper [12], and refrain here from repeating its details.

2.3 Solving PLDEs in $\Pi\Sigma^*$ -Extensions

In his seminal paper, Karr [10] has introduced the notion of $\Pi\Sigma$ -fields for representing nested sum and product expressions in difference fields. A simplified version of these fields are the $\Pi\Sigma^*$ -fields [20]. These are difference fields which are obtained from a constant field \mathbb{K} by adjoining formal sums or products. To be precise, $(\mathbb{G}(t), \sigma)$ is called a $\Pi\Sigma^*$ -extension of a difference field (\mathbb{G}, σ) , if both difference fields share the same field of constants, t is transcendental over \mathbb{G} , and $\sigma(t) = t + r$ for some $r \in \mathbb{G}$ (then t represents a sum) or $\sigma(t) = r \cdot t$ for some $r \in \mathbb{G}$ (then t represents a product). A $\Pi\Sigma^*$ -field is a difference field $(\mathbb{K}(t_1, \dots, t_e), \sigma)$ which is obtained from a constant field \mathbb{K} by repeated $\Pi\Sigma^*$ -extensions. In short, we say that $(\mathbb{G}(t_1, \dots, t_e), \sigma)$ is a (nested) $\Pi\Sigma^*$ -extension of (\mathbb{G}, σ) if all the t_i are $\Pi\Sigma^*$ -extensions.

It is known how to solve PLDEs in $\Pi\Sigma^*$ -fields [25]. Here, we are interested in difference fields which are obtained by building a tower of $\Pi\Sigma^*$ -extensions on top of a free difference field.

Example 2. For representing the sum from Example 1 above, we choose the difference field $(\mathbb{Q}\langle x \rangle(k), \sigma)$, where $(\mathbb{Q}\langle x \rangle, \sigma)$ is a free difference field and $\sigma(k) = k + 1$. In this field, $f_{n+i, k+j} = (k+j)X_{(n+i)+(k+j)}$ ($i, j \in \mathbb{Z}$) is represented by $(k+j)x_{i+j}$. The creative telescoping equation admits a solution (c_0, c_1, c_2, g) in the same field (compare Example 1). The recurrence obtained for the whole sum, however, requires besides n and X_n the representation of X_{2n} and X_{2n+1} . We take $(\mathbb{Q}\langle x \rangle \langle y \rangle \langle z \rangle(n), \sigma)$ with x, y, z free (representing X_n , X_{2n} , and X_{2n+1} , respectively) and $\sigma(n) = n + 1$. The solutions 1 and n of the homogeneous equation can be represented in the same field, but for representing a particular solution, we have to change to the bigger field $(\mathbb{Q}\langle x \rangle \langle y \rangle \langle z \rangle(n)(t_1)(t_2), \sigma)$, which is a $\Pi\Sigma^*$ -extension of the original one [3, 20]. Here $\sigma(t_1) = t_1 + \sigma^{-1}(f)$ and $\sigma(t_2) = t_2 + \sigma(t_1)$, where

$$f = -(-x_1 + (n+1)z_0 + (n+2)y_1 - (n+1)z_1 - (n+2)y_2).$$

A particular solution is then t_2 (compare again Example 1).

Subsequently, let \mathbb{K} be a field of constants, \mathbb{G} be obtained from \mathbb{K} by extension of zero or more free difference variables, and \mathbb{F} be obtained from \mathbb{G} by zero or more $\Pi\Sigma^*$ -extensions, say $\mathbb{F} = \mathbb{G}(t_1, \dots, t_e)$. In such a field, PLDEs can be solved

very much as in $\Pi\Sigma^*$ fields. Let $\mathbb{F}(t)$ be a $\Pi\Sigma^*$ -extension of \mathbb{F} . We outline a method which reduces the problem of solving a PLDE in $\mathbb{F}(t)$ to solving several PLDEs in \mathbb{F} . Only a rough overview is given, some remarks on technical details and pointers to literature are given in the end of the section.

Reduction I (denominator bounding). Compute a nonzero polynomial $d \in \mathbb{F}[t]$ such that for all $c_i \in \mathbb{K}$ and $g \in \mathbb{F}(t)$ with (5) we have $dg \in \mathbb{F}[t]$. Then it follows that

$$\frac{a_0}{d}g' + \dots + \frac{a_m}{\sigma^m(d)}\sigma^m(g') = c_0f_0 + \dots + c_rf_r \quad (6)$$

for $g' \in \mathbb{F}[t]$ if and only if (5) with $g = g'/d$.

Reduction II (degree bounding). Given such a denominator bound, it suffices to look only for $c_i \in \mathbb{K}$ and polynomial solutions $g \in \mathbb{F}[t]$ with (5). Next, we compute a degree bound $b \in \mathbb{N}_0$ for these polynomial solutions.

Reduction III (polynomial degree reduction). Given such a degree bound one looks for $c_i \in \mathbb{K}$ and $g_i \in \mathbb{F}$ such that (5) holds for $g = \sum_{i=0}^b g_i t^i$. This can be achieved as follows. First derive the possible leading coefficients g_b by solving a specific PLDE in (\mathbb{F}, σ) , then plug its solutions into (5) and recursively look for the remaining solutions $g = \sum_{i=0}^{b-1} g_i t^i$. Thus one can derive the solutions of a PLDE over $(\mathbb{F}(t), \sigma)$ by solving several PLDEs in (\mathbb{F}, σ) .

As already worked out in [12], this reduction leads us to a complete algorithm that solves PLDEs for $m = 1$. Moreover, in [25] based on results from [5, 21, 23, 12] we show that this reduction delivers a method that eventually produces all solutions for the higher order case $m \geq 2$.

Finally, we mention a refined version of parameterized telescoping (5) ($m = 1, a_0 = -1, a_1 = 1$). If no solution exists in \mathbb{F} , we can decide if there exists a solution in a $\Pi\Sigma^*$ -extension of \mathbb{F} where the sums and products are not more deeply nested than the original expressions in the f_i . If \mathbb{F} is a $\Pi\Sigma$ -field, this problem has been solved in [24]. Also this algorithm carries over if \mathbb{F} contains free variables; see Remark 4 for details.

Example 3. Applying our refined telescoping algorithm to the sum (4) gives the identity

$$\begin{aligned} \sum_{k=0}^n kX_{n+k} &= (n+1) \sum_{k=1}^n X_{k-1} - \sum_{k=1}^n kX_{k-1} \\ &\quad - (n+2) \sum_{k=1}^n X_{2k-2} + 2 \sum_{k=1}^n kX_{2k-2} \\ &\quad - (n+1) \sum_{k=1}^n X_{2k-1} + 2 \sum_{k=1}^n kX_{2k-1} + nX_{2n}. \end{aligned}$$

Despite being more lengthy, this representation is preferred, because it only contains indefinite sums.

Remark 1. (Denominator bounding) It was shown in [12] that a denominator bound d can be computed if $m = 1$. For the case $m \geq 2$ this problem is not completely solved: For Σ^* -extensions we still can compute a denominator bound, but if t is a Π -extension, we find the denominator bound only up to a power of t . The corresponding algorithms can be found in [21, Algorithm 1] by combining certain subproblems solved in [12, Corollary 1, Theorem 2, Theorem 4].

Remark 2. (Degree bounding) A degree bound can be computed if $m = 1$. For the case $m \geq 2$ we have algorithms only for various special cases [23].

Remark 3. (Degree reduction) Following the reduction recursively, one can solve PLDEs in $\mathbb{F}(t)$ if one can compute all the needed denominators (Reduction I) in $\mathbb{F}[t]$ and $\mathbb{G}(t_1) \dots (t_{i-1})[t_i]$, the degree bounds (Reduction II) in $\mathbb{F}[t]$ and $\mathbb{G}(t_1) \dots (t_{i-1})[t_i]$, and all the resulting PLDEs in \mathbb{G} obtained by recursive application of Reduction III. Here the following remarks are in place.

(1) If \mathbb{G} is the constant field, solving PLDEs in \mathbb{G} reduces to linear algebra [25, page 805]. Also if \mathbb{G} is a free difference field, PLDEs can be solved; see Section 2.2.

(2) If $m = 1$, degree and denominator bounds can be computed in $\mathbb{F}[t]$ and $\mathbb{G}(t_1) \dots (t_{i-1})[t_i]$; see Remarks 1 and 2. Hence we get a complete algorithm for solving PLDEs.

(3) If $m \geq 2$, denominator and degree bounds can be computed only partially so far. But, as worked out in [25, Theorem 5.3], the reduction leads to a recursive enumeration procedure that eventually outputs all solutions of a given PLDE in $\mathbb{F}(t)$.

Remark 4. (Refined telescoping) We obtain the algorithm for this problem by combining results from [21, 23, 24, 12]. Namely, there are algorithms for computing denominator and degree bounds which have the additional property that they are extension-stable; this follows by [21, Thm. 10], [23, Thm. 17], and the fact that one can handle certain subproblems in \mathbb{F} ; see [12, Theorems 1, 3, 4]. Using this fact, we obtain an algorithm that solves the refined telescoping problem; see [24, Thms. 6, 8].

3. NEGATIVE RESULTS

With an implementation of the algorithm described in the previous section, we have searched for variations of the definite sum $\sum_{k=0}^n kX_{n+k}$ of Example 1. These experiments have led us to the following theorem, which gives an explicit a priori criterion for which sequences $f_{n,k}$ a linear recurrence for the general definite sum $\sum_{k=0}^n f_{n,k}X_{n+k}$ is found.

THEOREM 1. *Let (\mathbb{F}, σ) be a difference field with constant field \mathbb{K} , and let $f_0, \dots, f_r \in \mathbb{F}$. Then there exist $c_0, \dots, c_r \in \mathbb{K}$ and $g \in \mathbb{F}(x)$ with*

$$\sigma(g) - g = c_0 f_0 x_0 + c_1 f_1 x_1 + \dots + c_r f_r x_r$$

if and only if there exist $b_0, \dots, b_r \in \mathbb{K}$ with

$$b_0 \sigma^r(f_0) + b_1 \sigma^{r-1}(f_1) + \dots + b_r f_r = 0.$$

PROOF. Suppose first that f_0, \dots, f_r are such that there exist $b_0, \dots, b_r \in \mathbb{K}$ with

$$b_0 \sigma^r(f_0) + b_1 \sigma^{r-1}(f_1) + \dots + b_r f_r = 0. \quad (7)$$

Then $c_k := b_k$ ($k = 0, \dots, r$) and

$$g := a_0 x_0 + a_1 x_1 + \dots + a_{r-1} x_{r-1},$$

where $a_k = -\sum_{i=0}^k b_i \sigma^{k-i}(f_i)$ are as required: We have

$$\begin{aligned} \sigma(g) - g &= -a_0 x_0 + \sum_{k=1}^{r-1} (\sigma(a_{k-1}) - a_k) x_k + \sigma(a_{r-1}) x_r \\ &= b_0 f_0 x_0 + \dots + b_r f_r x_r, \end{aligned}$$

because

$$\begin{aligned} -a_0 &= b_0 f_0 \\ \sigma(a_{k-1}) - a_k &= -\sum_{i=0}^{k-1} b_i \sigma^{(k-1)-i+1}(f_i) + \sum_{i=0}^k b_i \sigma^{k-i}(f_i) \\ &= b_k f_k \quad (k = 1, \dots, r-1) \\ \sigma(a_{r-1}) &= -\sum_{i=0}^{r-1} b_i \sigma^{r-i}(f_i) \stackrel{(7)}{=} b_r f_r. \end{aligned}$$

This proves the first implication. Now, assume that there exist $c_0, \dots, c_r \in \mathbb{K}$ and $g \in \mathbb{F}(x)$ such that

$$\sigma(g) - g = c_0 f_0 x_0 + c_1 f_1 x_1 + \dots + c_r f_r x_r. \quad (8)$$

The element g is a rational function in x_i ($i \in \mathbb{Z}$) with coefficients in \mathbb{F} . However, g can have neither a nontrivial denominator nor a nonlinear term w.r.t. the x_i , because then $\sigma(g) - g$ would have nontrivial denominator or a nonlinear term as well. Furthermore, g must be free of all x_i with $i < 0$ or $i \geq r$, for otherwise $\sigma(g) - g$ would contain some x_i with $i < 0$ or $i > r$ in mismatch with the right hand side of (8). Thus g can only have the form

$$g = a_0 x_0 + a_1 x_1 + \dots + a_{r-1} x_{r-1}$$

for certain $a_i \in \mathbb{F}$. Now

$$\sigma(g) - g = -a_0 x_0 + \sum_{k=1}^{r-1} (\sigma(a_{k-1}) - a_k) x_k + \sigma(a_{r-1}) x_r,$$

and comparing coefficients of x_k ($k = 0, \dots, r-1$) with the right hand side of (8) gives

$$\begin{aligned} a_0 &= -c_0 f_0 \\ a_1 &= \sigma(a_0) - c_1 f_1 = -(c_0 \sigma(f_0) + c_1 f_1) \\ &\vdots \end{aligned}$$

$$a_{n-1} = \sigma(a_{r-2}) - c_{r-1} f_{r-1} = -\sum_{i=0}^{r-1} c_i \sigma^{(r-1)-i}(f_i).$$

Comparing finally the coefficient of x_r gives

$$c_r f_r = \sigma(a_{r-1}) = -\sum_{i=0}^{r-1} c_i \sigma^{(r-1)-i+1}(f_i),$$

and therefore we must have

$$c_0 \sigma^r(f_0) + c_1 \sigma^{r-1}(f_1) + \dots + c_r f_r = 0,$$

as claimed. \square

For obtaining a recurrence equation of the definite sum $S_n := \sum_{k=0}^n f_{n,k} X_{n+k}$ via creative telescoping, we choose $f_0 = f_{n,k}$, $f_1 = f_{n+1,k}$, \dots , $f_r = f_{n+r,k}$ (or more precisely, corresponding difference field elements). Thus the above theorem states that a recurrence for S_n is found via creative telescoping if and only if the antidiagonal sequences $f_{i-k,k}$ ($i \in \mathbb{Z}$) are all solutions of a single homogeneous linear recurrence whose coefficients are constant with respect to k :

COROLLARY 1. *Creative telescoping delivers a recurrence for a definite sum $\sum_{k=0}^n f_{n,k} X_{n+k}$, where $f_{n,k}$ is free of X_k , if and only if there exist constants b_0, \dots, b_r such that*

$$b_0 f_{n,k+r} + b_1 f_{n+1,k+r-1} + \dots + b_r f_{n+r,k} = 0$$

for all n and k . \square

Similar criteria can be obtained for sums of the form

$$S_n = \sum_{k=0}^n f_{n,k} X_{an+bk}$$

for any fixed $a, b \in \mathbb{Z}$. For $a = 1, b = -1$, we find the criterion that $c_0, \dots, c_r \in \mathbb{K}$ and $g \in \mathbb{F}\langle x \rangle$ with

$$\sigma(g) - g = c_0 f_0 x_0 + c_1 f_1 x_{-1} + \dots + c_r f_r x_{-r}$$

exist iff there exist $b_0, \dots, b_r \in \mathbb{K}$ with

$$b_0 f_0 + b_1 \sigma(f_1) + \dots + c_r \sigma^r(f_r) = 0,$$

If $\gcd(a, b) = 1$ (which we may assume without loss of generality by the substitution $X'_i := X_{\gcd(a,b)i}$) and $|a| > 1$, then a recurrence exists only in the trivial case $f_{n,k} \equiv 0$. The case $|a| = 1, |b| \geq 0$ leads to a restriction on the summand similar as the one stated above. The arguments for all these variations are fully analogous to the proof given above.

The theorem provides a means to obtain creative telescoping recurrences without actually executing the algorithm described in Section 2.1.

Example 4. Consider once more the definite sum $S_n = \sum_{k=0}^n k X_{n+k}$ of Example 1. In the notation of the theorem, we have $f_{n,k} = k$, so we immediately obtain the recurrence

$$g_{n,k+1} - g_{n,k} = k X_{n+k} - 2k X_{(n+1)+k} + k X_{(n+2)+k},$$

where

$$g_{n,k} = (k-1) X_{n+k+1} - k X_{n+k}.$$

Summing on k from 0 to $n+2$ leads to the same inhomogeneous recurrence which we obtained before.

The more interesting implication of Theorem 1 and Corollary 1 is of course that which allows us to definitely exclude the existence of creative telescoping recurrences for sums of a certain shape. For instance, the following results follow immediately.

COROLLARY 2. *For the following definite sums, no recurrence can be found via creative telescoping.*

1. $\sum_{k=0}^n r(k) X_{n+k}$ for any $r \in \mathbb{K}(k) \setminus \mathbb{K}[k]$,
2. $\sum_{k=0}^n \binom{n}{k} X_{n+k}$,
3. $\sum_{k=0}^n r(k) \prod_{i=1}^k X_{n+i}$ for any $r \in \mathbb{K}(k) \setminus \mathbb{K}[k]$. \square

For the first two sums, the statement follows from the fact that rational functions and the binomial coefficients do not form solutions of homogeneous recurrence equations with constant coefficients. The third sum can be brought to the form of Corollary 1 by putting $Y_k := \prod_{i=0}^k X_i$, for then

$$\sum_{k=0}^n r(k) \prod_{i=1}^k X_{n+i} = \sum_{k=0}^n r(k) Y_{n+k} / Y_n = \frac{1}{Y_n} \sum_{k=0}^n r(k) Y_{n+k},$$

and the factor $1/Y_n$, which is independent of k , does not affect the existence of a recurrence. The possibility of such substitutions extends the range of Corollary 1. In order to find nontrivial examples, it is necessary to focus on sums with more complicated summands.

4. POSITIVE RESULTS

We present in this section some general identities which we have found by using the algorithm described in Section 2.1. To our knowledge, most of these identities are new. The examples are separated into indefinite and definite sums.

4.1 Indefinite summation

Example 5. For the sum $\sum_{k=1}^a (-1)^k \sum_{j=1}^k X_j$ we find the representation

$$\sum_{k=1}^a (-1)^k \sum_{j=1}^k X_j = \frac{1}{2} \left[(-1)^a \sum_{k=1}^a X_k + \sum_{k=1}^a (-1)^k X_k \right].$$

Specializing X_j gives the following identities.

- $X_j := \frac{1}{j}$:

$$\sum_{k=1}^a (-1)^k H_k = \frac{1}{2} (-1)^a H_a + \frac{1}{2} \sum_{k=1}^a \frac{(-1)^k}{k};$$

$H_a = \sum_{k=1}^a \frac{1}{k}$ denotes the harmonic numbers.

- $X_j := \binom{n}{j-1}, a := n+1$; see [27, Thm. 4.2]:

$$\sum_{k=0}^n (-1)^{k+1} \sum_{j=0}^k \binom{n}{j} = \frac{1}{2} (-1)^{n+1} 2^n.$$

Example 6. We find

$$\sum_{k=1}^a (-1)^k \binom{n}{k} \sum_{j=1}^k X_j = \frac{1}{n} \left[(n-a) \binom{n}{a} (-1)^a \sum_{k=1}^a X_k + \sum_{k=1}^a (-1)^k k \binom{n}{k} X_k \right].$$

In particular, in the special case $a = n$ and $X_j = \frac{1}{j^m}$ where $m \geq 1$ we obtain the following simplification. By [13, Prop. 2.1] we get

$$\sum_{k=1}^n \frac{(-1)^k}{k^{m-1}} \binom{n}{k} = -\frac{1}{(m-1)!} B_{m-1}(\dots, (i-1)! H_N^{(i)}, \dots)$$

where $B_m(\dots, x_i, \dots)$ are the complete Bell polynomials [7] and

$$H_n^{(s)} = \sum_{k=1}^n \frac{1}{k^s}, \quad s > 0$$

are the generalized harmonic numbers. Hence we arrived at the representation

$$\sum_{k=1}^n (-1)^k \binom{n}{k} H_k^{(m)} = -\frac{1}{n(m-1)!} B_{m-1}(\dots, (i-1)! H_n^{(i)}, \dots);$$

see [15, Thm. 3]. The first instances are:

$$\sum_{k=1}^a (-1)^k \binom{n}{k} H_k = -\frac{1}{n},$$

$$\sum_{k=1}^a (-1)^k \binom{n}{k} H_k^{(2)} = -\frac{1}{n} H_n,$$

$$\sum_{k=1}^a (-1)^k \binom{n}{k} H_k^{(3)} = -\frac{1}{2n} (H_n^2 + H_n^{(2)}).$$

Further indefinite summation identities are

$$\begin{aligned} & \sum_{k=1}^a (-1)^k \left(\sum_{j=1}^k X_j - \frac{X_k}{2} \right)^2 \\ &= \frac{1}{2} (-1)^a \left(\sum_{k=1}^a X_k \right)^2 - \frac{1}{4} \sum_{k=1}^a (-1)^k X_k^2, \\ & \sum_{k=1}^a \left(\sum_{j=1}^k X_j + X_k(k-1) \right)^2 \\ &= a \left(\sum_{k=1}^a X_k \right)^2 - \sum_{k=1}^a k X_k^2 + \sum_{k=1}^a k^2 X_k^2. \end{aligned}$$

4.2 Definite summation and summability criteria

Example 7. Similarly as in Examples 1 and 3 we find (and prove) with our difference field machinery the identity

$$\sum_{k=0}^n \left(\sum_{i=0}^k X_{n-i} \right)^2 = 2 \sum_{k=0}^n X_k \sum_{j=0}^k j X_{j-1} + \sum_{k=0}^n X_k^2 + \sum_{k=0}^n k X_k^2.$$

Namely, starting with $S_n = \sum_{k=0}^n \left(\sum_{i=0}^k X_{n-i} \right)^2$ we can compute by creative telescoping the recurrence

$$\begin{aligned} & -X_{n+1} S_{n+2} + (X_{n+1} + X_{n+2}) S_{n+1} - X_{n+2} S_n \\ &= -X_{n+1} X_{n+2} ((n+2) X_{n+1} + (n+3) X_{n+2}). \end{aligned}$$

Next, we solve this recurrence relation and find the solutions $1, \sum_{k=0}^n X_k$ for the homogeneous version and the particular solution

$$P_n = \sum_{k=0}^n X_k \sum_{i=0}^k (i X_{i-1} + X_i + i X_i);$$

since $S_n = P_n$ for $n = 0, 1$, it follows that $P_n = S_n$ for all n . Finally, applying our indefinite summation algorithm, we get the simplification

$$S_n = P_n = 2 \sum_{k=0}^n X_k \sum_{j=0}^k j X_{j-1} + \sum_{k=0}^n X_k^2 + \sum_{k=0}^n k X_k^2.$$

Example 8. Consider the definite sum

$$S_n := \sum_{k=1}^n \binom{n}{k} \sum_{j=1}^k X_j.$$

With our refined creative telescoping algorithm we can compute the recurrence

$$S_{n+1} - 2S_n = \sum_{k=0}^n \binom{n}{k} X_{k+1}. \quad (9)$$

In the next step we would like to solve this recurrence. For applying our algorithms, we first have to express the definite sum

$$C_n := \sum_{k=0}^n \binom{n}{k} X_{k+1}$$

in terms of $\Pi\Sigma^*$ -expressions. To this end, we would normally compute a recurrence for that sum and afterwards solve it. However, no recurrence can be found in this case.

(This can also be seen with Corollary 1, because $C_n = \sum_{k=0}^n \binom{n}{k} X_{n+1-k}$ after reversing the order of summation). Therefore, we try to specialize X_k in such a way that C_n can be written in a simple form (e.g., can be represented in form of a $\Pi\Sigma^*$ -extension) and that the solutions of (9) are nice. Note that (9) gives us a recipe for discovering nice identities. We may therefore consider (9) as a *summability criterion* for the sum S_n .

• $X_k = \frac{1}{k}$: We get easily the identity $C_n = \frac{2^{n+1}-1}{n+1}$ and find the general solution

$$c2^n + 2^n \left(H_n - \sum_{k=1}^n \frac{1}{k2^k} \right)$$

for (9). By choosing $c = 0$ we obtain

$$\sum_{k=1}^n \binom{n}{k} H_k = 2^n \left(H_n - \sum_{k=1}^n \frac{1}{k2^k} \right);$$

see [17, Equ. (41)].

With the same strategy we get the following identities.

• $X_k = \frac{1}{k^2}$:

$$\sum_{k=1}^n \binom{n}{k} H_k^{(2)} = 2^n \left(\sum_{k=1}^n \frac{\sum_{j=1}^k \frac{2^j}{j}}{k2^k} - \sum_{k=1}^n \frac{H_k}{k2^k} \right).$$

• $X_k = (k-1)!$:

$$\sum_{k=1}^n \binom{n}{k} \sum_{j=1}^k (j-1)! = 2^n \left(\sum_{k=1}^n \frac{k!}{k2^k} + \sum_{k=1}^n \frac{k! \sum_{j=1}^k \frac{1}{j!}}{k2^k} - \sum_{k=1}^n \frac{1}{k2^k} \right).$$

Example 9. We attack the sum

$$S_n := \sum_{k=1}^n \binom{n}{k}^2 \sum_{j=1}^k X_j$$

as in the previous example. First, we compute the recurrence

$$(n+1)S_{n+1} - 2(2n+1)S_n = \sum_{k=0}^n (3n-2k+1) \binom{n}{k}^2 X_{k+1}.$$

Looking at this summability criterion we find the following identities.

• $X_k = \frac{1}{k}$; see [8, Equ. 2.26]:

$$\sum_{k=1}^n \binom{n}{k}^2 H_n = \binom{2n}{n} (2H_n - H_{2n}).$$

• $X_k = \frac{1}{k^2}$:

$$\sum_{k=1}^n \binom{n}{k}^2 H_k^{(2)} = \binom{2n}{n} (2H_n^{(2)} - 3 \sum_{k=1}^n \frac{1}{k^2 \binom{2k}{k}}).$$

Example 10. We compute for

$$S_n = \sum_{k=0}^n X_k \sum_{i=1}^{n-k} Y_i$$

the recurrence

$$S_{n+1} - S_n = \sum_{k=0}^n X_k Y_{n-k+1}.$$

With this summability criterion we find the following identities.

- $X_k = \frac{1}{k!}, Y_k = \frac{1}{(k-2)!}$:

$$\sum_{k=0}^n \frac{1}{k!} \sum_{j=1}^{n-k} \frac{1}{(j-2)!} = \frac{1}{2} \sum_{k=1}^{n-1} \frac{2^k}{(k-1)!}.$$

- $X_k = H_k, Y_k = \frac{1}{k}$; see [14, Chapter 1.2.7, Exercise 22]:

$$\sum_{k=0}^n H_k H_{n-k} = 2n(1 - H_n) + (n+1)H_n^2 - (n+1)H_n^{(2)}.$$

5. CONCLUSION

The extension of symbolic summation algorithms to free difference fields allows one the discovery of general families of summation identities, depending on unspecified sequences. We have illustrated in this paper how the $\Pi\Sigma$ -theory for nested sum expressions can be extended, and we have found several general identities with the modified summation algorithms. We have also indicated (Section 3) that interesting relations can only be found for sums whose summand exceeds a certain level of sophistication; if it is too simple, then only trivial relations remain.

Though our extension itself is not very difficult, it should be remarked that it is based on a highly developed machinery for generating and solving recurrence equations with difference fields. Without using, for instance, algorithms that can optimize the nesting depth of sum expressions, we would hardly have been able to find any of the nontrivial examples in Section 4. Once general identities like those of Section 4 are available, they may be specialized in such a way that well-developed theories and/or algorithms can be applied for further processing. Example 6 points into that direction.

Acknowledgement. We would like to thank the three anonymous referees for their valuable remarks.

6. REFERENCES

- [1] S. Abramov, J. Carette, K. Geddes, and H. Le. Telescoping in the context of symbolic summation in Maple. *Journal of Symbolic Computation*, 38(4):1303–1326, 2004.
- [2] S. A. Abramov. A criterion for validity of the discrete Newton-Leibniz formula. In *this volume*, 2006.
- [3] S. A. Abramov and M. Petkovšek. D’Alembertian solutions of linear differential and difference equations. In J. von zur Gathen, editor, *Proc. ISSAC’94*, pages 169–174. ACM Press, Baltimore, 1994.
- [4] S. A. Abramov and M. Petkovšek. Gosper’s algorithm, accurate summation, and the discrete Newton-Leibniz formula. In *Proceedings of ISSAC’06*, pages 5–12, 2005.
- [5] M. Bronstein. On solutions of linear ordinary difference equations in their coefficient field. *J. Symbolic Comput.*, 29(6):841–877, June 2000.
- [6] R. M. Cohn. *Difference Algebra*. Interscience Publishers, John Wiley & Sons, 1965.
- [7] L. Comtet. *Advanced Combinatorics*. D. Reidel Publishing Company, 1970.
- [8] G. P. Egorychev. *Integral Representation and the Computation of Combinatorial Sums*. Translations of Mathematical Monographs. American Mathematical Society, Providence, RI, 1984. Translated from the Russian by H. H. McFadden. Translation edited by Lev J. Leifman.
- [9] W. Gosper. Decision procedure for indefinite hypergeometric summation. *Proceedings of the National Academy of Sciences of the United States of America*, 75:40–42, 1978.
- [10] M. Karr. Summation in finite terms. *J. ACM*, 28:305–350, 1981.
- [11] M. Kauers. Computer proofs for polynomial identities in arbitrary many variables. In *Proceedings of ISSAC’04*, pages 199–204, July 2004.
- [12] M. Kauers and C. Schneider. Indefinite summation with unspecified sequences. *Discrete Mathematics*, to appear.
- [13] P. Kirschenhofer. A note on alternating sums. *Electron. J. Combin.*, 3(2), 1996.
- [14] D. Knuth. *The art of computer programming*, volume 1, Fundamental Algorithms. Addison Wesley, Reading, MA, 3rd edition, 2000.
- [15] P. Larcombe, M. Larsen, and E. Fennessey. On two classes of identities involving harmonic numbers. *Util. Math.*, 67:65–80, 2005.
- [16] P. Paule. Greatest factorial factorization and symbolic summation. *J. Symbolic Comput.*, 20:235–268, 1995.
- [17] P. Paule and C. Schneider. Computer proofs of a new family of harmonic number identities. *Adv. in Appl. Math.*, 31(2):359–378, 2003.
- [18] P. Paule and M. Schorn. A Mathematica version of Zeilberger’s algorithm for proving binomial coefficient identities. *Journal of Symbolic Computation*, 20(5–6):673–698, 1995.
- [19] M. Petkovšek, H. S. Wilf, and D. Zeilberger. *A = B*. A. K. Peters, Wellesley, MA, 1996.
- [20] C. Schneider. Symbolic summation in difference fields. Technical Report 01-17, RISC-Linz, J. Kepler University, 2001. PhD Thesis.
- [21] C. Schneider. A collection of denominator bounds to solve parameterized linear difference equations in $\Pi\Sigma$ -extensions. In D. P. et.al., editor, *Proc. SYNASC04, 6th Internat. Symposium on Symbolic and Numeric Algorithms for Scientific Computation*, pages 269–282. Mirton Publishing, 2004.
- [22] C. Schneider. The summation package Sigma: Underlying principles and a rhombus tiling application. *Discrete Math. Theor. Comput. Sci.*, 6(2):365–386, 2004.
- [23] C. Schneider. Degree bounds to find polynomial solutions of parameterized linear difference equations in $\Pi\Sigma$ -fields. *Appl. Algebra Engrg. Comm. Comput.*, 16(1):1–32, 2005.
- [24] C. Schneider. Finding telescopers with minimal depth for indefinite nested sum and product expressions. In M. Kauers, editor, *Proc. ISSAC’05*, pages 285–292. ACM, 2005.
- [25] C. Schneider. Solving parameterized linear difference equations in terms of indefinite nested sums and products. *J. Differ. Equations Appl.*, 11(9):799–821, 2005.
- [26] D. Zeilberger. The method of creative telescoping. *J. Symbolic Comput.*, 11:195–204, 1991.
- [27] Z. Zhang. A kind of binomial identity. *Discrete Math.*, 196:291–298, 1999.