

COMPUTER ALGEBRA AND POWER SERIES WITH POSITIVE COEFFICIENTS

MANUEL KAUERS*

ABSTRACT. We consider the question whether all the coefficients in the series expansions of some specific rational functions are positive, and we demonstrate how computer algebra can help answering questions arising in this context. By giving partial computer proofs, we provide new evidence in support of some longstanding open conjectures. Also two new conjectures are made.

AMS SUBJECT CLASSIFICATION. 33F10, 13F25, 05E99

KEYWORDS. Computer Algebra, Positivity Theory, Power Series.

1. INTRODUCTION

Proving that all the coefficients in the series expansion of some given multivariate rational function are positive can be quite a difficult task. There are difficult papers on this subject by Szegő [14], Askey and Gasper [2], Koornwinder [10], and others. Gillis, Reznick and Zeilberger [8] have pointed out that some seemingly difficult positivity results can be proven also by elementary means. In this paper, we make an attempt at going one step further: We ask to which extent positivity results can be proven automatically using computer algebra. Two results from the literature and two longstanding open conjectures related to them are considered. For none of the latter, we are able to provide full proofs, but we give partial proofs that add new evidence in support of these conjectures.

Also Zeilberger [15] addresses the question of proving positivity results with the aid of computer algebra. He proposes a method using positivity-preserving transformations, which is independent of our approach described below. The approach we take is a continuation of our previous work on treating special function inequalities via symbolic computation [5, 6, 9, 1].

While it is easy to show that there can be no algorithm which for a given multivariate rational function decides whether all its series coefficients are positive, computer algebra is nevertheless useful for deciding subproblems that may arise in the construction of a positivity proof. Our proofs follow a common pattern: We first determine recurrence equations for the coefficients (using computer algebra, see Section 1.1), and then prove a suitable quantified formula about polynomial inequalities (using computer algebra, see Section 1.2) which together with the recurrence equations implies the desired positivity result. Sometimes this method is successful, sometimes it is not.

Before entering the subject, let us briefly summarize the two main techniques from computer algebra that are used throughout the rest of the paper.

1.1. Guessing and Proving Recurrence Relations. Let $a(n, m)$ be a sequence whose value can be computed for every particular point $(n, m) \in \mathbb{N}^2$. If the sequence satisfies a recurrence equation with polynomial coefficients, then this equation can be found easily by making an ansatz. To this end, we first choose a finite set $S \subseteq \mathbb{N}^4$. A recurrence corresponds to an array of constants α_i ($i \in S$) with

$$\sum_{i=(i_1, i_2, i_3, i_4) \in S} \alpha_i n^{i_1} m^{i_2} a(n + i_3, m + i_4) = 0.$$

In order to find candidates for α_i , we evaluate the above expression for a couple of specific sample points $(n, m) \in \mathbb{N}^2$ and undetermined coefficients α_i , thus obtaining a linear system for the

* Partially supported by the Austrian Science Foundation (FWF) grants SFB F1305 and P16613-N12.

coefficients α_i . Each coefficient vector $(\alpha_i : i \in S)$ corresponding to an actual recurrence will belong to the solution space of the system, but there might be additional solutions not corresponding to recurrence equations. This is why the method is referred to as *automated guessing* [12].

If the $a(n, m)$ arise as coefficients in the series expansion of a rational function $r(x, y)$, then it is easy to decide for a conjectured recurrence whether it is true or false. It suffices to transform the conjectured recurrence into a PDE for its generating function (this can be done automatically) and check whether $r(x, y)$ solves that differential equation. It will do so if and only if the conjectured recurrence holds true. We can thus repeat the guessing procedure with bigger and bigger sets of sample points until the recurrences delivered by the guessing method are all found to be true.

It is of course immaterial that $a(n, m)$ is a bivariate sequence; the same algorithm is applicable for any arity. All the recurrence equations claimed in this paper have been found and verified in this way, unless otherwise stated.

1.2. Proving and Finding Polynomial Inequalities. Our second ingredient is the CAD algorithm [4]. This algorithm operates on quantified formulas about polynomial inequalities over the reals. Formally, a quantifier-free formula $\Phi(x_1, \dots, x_n)$ is

- a logical constant, i.e., True or False,
- or an atomic formula, i.e., $p(x_1, \dots, x_n) \diamond 0$ for a polynomial $p \in k[x_1, \dots, x_n]$ and a relation $\diamond \in \{=, \neq, >, <, \geq, \leq\}$; k being the field of real algebraic numbers,
- or a boolean combination of other quantifier-free formulas, i.e.,

$$\Psi_1(x_1, \dots, x_n) \diamond \Psi_2(x_1, \dots, x_n)$$

with $\diamond \in \{\wedge, \vee, \Rightarrow, \Leftrightarrow\}$, or $\neg\Psi_1(x_1, \dots, x_n)$, for some subformulas Ψ_1, Ψ_2 .

A quantified formula $\Phi(x_1, \dots, x_n)$ is a formula of the form

$$Q_1y_1 Q_2y_2 \dots Q_my_m : \Psi(x_1, \dots, x_n, y_1, \dots, y_m),$$

where Ψ is a quantifier-free formula and $Q_1, \dots, Q_m \in \{\forall, \exists\}$. The x_1, \dots, x_n are referred to as free variables, as opposed to the y_1, \dots, y_m which are called bounded variables.

The CAD algorithm is able to perform quantifier elimination over such formulas, i.e., given any quantified formula $\Phi(x_1, \dots, x_n)$, it computes a quantifier-free formula $\Phi'(x_1, \dots, x_n)$ such that

$$\forall x_1, x_2, \dots, x_n \in \mathbb{R} : (\Phi(x_1, \dots, x_n) \iff \Phi'(x_1, \dots, x_n)).$$

The formula $\Phi'(x_1, \dots, x_n)$ describes the condition that the x_1, \dots, x_n have to satisfy in order to make Φ true. If CAD is applied to a formula Φ with no free variables, it will deliver as quantifier-free formula Φ' either True (then this is a rigorous proof for Φ) or False (then this is a rigorous proof for $\neg\Phi$).

All the quantified formulas claimed in this paper have been proven with Mathematica's implementation of CAD [13], unless otherwise stated.

2. A RESULT OF ASKEY AND GASPER, AND A CONJECTURED VARIATION

Theorem 1. [3] *Let $a(n, m, k)$ be such that*

$$\frac{1}{1 - (x + y + z) + 4xyz} = \sum_{n, m, k \geq 0} a(n, m, k) x^n y^m z^k.$$

Then $a(n, m, k) > 0$ for all $n, m, k \geq 0$.

First we determine all constants $\alpha_{i,u,v,w}$ such that

$$\sum_{u, v, w \in \{0, 1\}} (\alpha_{0, u, v, w} + \alpha_{1, u, v, w} n + \alpha_{2, u, v, w} m + \alpha_{3, u, v, w} k) a(n + u, m + v, k + w) = 0$$

for all $n, m, k \geq 0$. It turns out that the possible choices for $\alpha_{i,u,v,w}$ form a vector space of dimension 10; the algorithm described in the introduction gives us a basis for this vector space, say the basis elements correspond to the recurrence equations

$$(1) \quad \sum_{u,v,w \in \{0,1\}} p_{i,u,v,w}(n, m, k) a(n+u, m+v, k+w) = 0 \quad (i = 1, \dots, 10),$$

for certain linear polynomials $p_{i,u,v,w}(n, m, k)$ that we know explicitly.

We want to construct a recurrence equation

$$\sum_{u,v,w \in \{0,1\}} q_{u,v,w}(n, m, k) a(n+u, m+v, k+w) = 0$$

with $q_{1,1,1}(n, m, k) < 0$ and $q_{u,v,w}(n, m, k) \geq 0$ ($(u, v, w) \neq (1, 1, 1)$) for all $n \geq m \geq k \geq 0$. From such a recurrence, the positivity of the $a(n, m, k)$ easily follows by induction. (It is sufficient to consider $n \geq m \geq k \geq 0$ by the symmetry of $a(n, m, k)$.) If such a recurrence exists, then it must be a linear combination of the basis equations in (1), so we can make an ansatz with undetermined coefficients:

$$(2) \quad q_{u,v,w}(n, m, k) := \beta_1 p_{1,u,v,w}(n, m, k) + \beta_2 p_{2,u,v,w}(n, m, k) + \dots + \beta_{10} p_{10,u,v,w}(n, m, k).$$

Next, we apply CAD to the formula

$$\forall n \geq m \geq k \geq 0 : q_{1,1,1}(n, m, k) < 0 \wedge \bigwedge_{\substack{u,v,w \in \{0,1\} \\ (u,v,w) \neq (1,1,1)}} q_{u,v,w}(n, m, k) \geq 0.$$

The resulting condition depends on the basis elements (1). For our basis, we obtained that the formula becomes true if and only if the β_i are chosen such that

$$\begin{aligned} & \beta_1 \leq 0 \wedge \beta_2 \geq 0 \wedge \beta_3 \geq 0 \wedge \beta_4 = 0 \wedge \beta_5 \geq 0 \wedge \beta_6 \geq 0 \wedge 2\beta_1 + 2\beta_2 + 2\beta_3 + \beta_4 + \beta_5 + \beta_6 \leq 0 \\ & \wedge \beta_3 \geq \beta_7 \wedge 2\beta_1 + \beta_3 + \beta_6 + \beta_7 \leq 0 \wedge \beta_2 + \beta_8 = 0 \wedge \beta_3 + \beta_4 \geq 2\beta_7 + \beta_8 + \beta_9 \\ & \wedge \beta_9 \leq 0 \wedge 3\beta_1 + \beta_2 + \beta_3 + \beta_5 + \beta_6 + \beta_7 + \beta_9 \leq 0 \wedge 2\beta_1 + \beta_5 + \beta_6 + 2(\beta_7 + \beta_9) \geq \beta_4 \\ & \wedge \beta_4 \leq \beta_5 + \beta_6 + 2\beta_{10} \wedge \beta_1 + \beta_3 + \beta_4 \geq \beta_7 + \beta_8 + \beta_{10} \wedge \beta_1 + \beta_2 + \beta_7 + \beta_8 \geq \beta_{10} \\ & \wedge \beta_{10} < 0 \wedge 2\beta_1 + \beta_2 + \beta_3 + \beta_5 + \beta_6 + \beta_{10} \leq 0 \wedge (\beta_1 + \beta_7 = 0 \vee (\beta_1 + \beta_7 < 0 \wedge \beta_1 + \beta_7 + \beta_9 \leq 0)). \end{aligned}$$

It is an easy matter to find a tuple $(\beta_1, \dots, \beta_{10})$ that satisfies this condition, if desired, we can have the computer find one. For instance, $(\beta_1, \dots, \beta_{10}) = (1, 0, 0, 0, 0, 2, 0, 0, 0, -1)$ satisfies the condition, and, if plugged into the ansatz equation (2), yields

$$2(n+m-k)a(n, m, k+1) + (1+n-m+k)a(n, m+1, k+1) - (1+n)a(n+1, m+1, k+1) = 0.$$

Together with the initial value $a(0, 0, 0) = 1 > 0$, this recurrence forms an easy induction proof of Theorem 1.

The recurrence above was first observed by Gillis and Kleeman [7]. They do, however, not remark on how they discovered this recurrence. As we have shown above, it is possible to derive it in a systematic way using computer algebra.

Let us now turn to an open problem. Gillis, Reznick and Zeilberger [8] have raised the conjecture that for any $r \geq 4$, the series coefficients of $1/(1 - (x_1 + x_2 + \dots + x_r) + r!x_1x_2 \dots x_r)$ are nonnegative. According to their Proposition 3, in order to confirm the conjecture, it suffices to prove nonnegativity of the diagonal coefficients $a_r(n, n, \dots, n)$ which are given by the binomial sum

$$a_r(n, n, \dots, n) = \sum_{k=0}^n (-1)^k \frac{(rn - (r-1)k)!(r!)^k}{(n-k)!r^k k!}.$$

As computational evidence for their conjecture, they verified that $a_r(n, n, \dots, n) \geq 0$ for $r = 4$ and $0 \leq n \leq 220$. We next prove the conjecture for $r = 4, 5, 6$ and arbitrary n .

Theorem 2.

(1) Let $a(n, m, k, l)$ be such that

$$\frac{1}{1 - (x + y + z + w) + 4!xyzw} = \sum_{n, m, k, l \geq 0} a(n, m, k, l) x^n y^m z^k w^l.$$

Then $a(n, m, k, l) \geq 0$ for all $n, m, k, l \geq 0$.

(2) Let $a(n, m, k, i, j)$ be such that

$$\frac{1}{1 - (x + y + z + u + v) + 5!xyzuv} = \sum_{n, m, k, i, j \geq 0} a(n, m, k, i, j) x^n y^m z^k u^i v^j.$$

Then $a(n, m, k, i, j) \geq 0$ for all $n, m, k, i, j \geq 0$.

(3) Let $a(n, m, k, i, j, l)$ be such that

$$\frac{1}{1 - (x + y + z + u + v + w) + 6!xyzuvw} = \sum_{n, m, k, i, j \geq 0} a(n, m, k, i, j) x^n y^m z^k u^i v^j w^l.$$

Then $a(n, m, k, i, j, l) \geq 0$ for all $n, m, k, i, j, l \geq 0$.

Proof. We prove the first statement, the others can be done in the same manner.

According to Prop. 3 of [8], it suffices to show that $a(n) \geq 0$ for all $n \geq 0$, where

$$a(n) = \sum_{k=0}^n (-1)^k \frac{(4n - 3k)! 4!^k}{(n - k)! 4! k!}.$$

With Zeilberger's algorithm [16, 11] we obtained the recurrence equation

$$\begin{aligned} & 331776(2n + 7)(4n + 11)(4n + 15)(n + 1)^3 a(n) \\ & + 13824(4n + 15) (32n^5 + 344n^4 + 1424n^3 + 2855n^2 + 2801n + 1085) a(n + 1) \\ & + 576 (192n^6 + 3072n^5 + 20108n^4 + 68918n^3 + 130513n^2 + 129613n + 52815) a(n + 2) \\ & - 8(n + 3)(4n + 7)(4n + 13) (40n^3 + 380n^2 + 1193n + 1240) a(n + 3) \\ & + (n + 4)^3(2n + 5)(4n + 7)(4n + 11)a(n + 4) = 0 \end{aligned}$$

for $a(n)$. Consider the formula

$$\forall A_0, A_1, A_2, A_3, A_4 \in \mathbb{R} \forall n \geq 0 :$$

$$(A_3 \geq \beta A_2 \wedge A_2 \geq \beta A_1 \wedge A_1 \geq \beta A_0 \geq 0 \wedge p_0(n)A_0 + \dots + p_4(n)A_4 = 0) \implies A_4 \geq \beta A_3,$$

where $p_i(n)$ denotes the polynomial appearing as coefficient of $a(n + i)$ in the above recurrence ($i = 0, \dots, 4$). Using CAD, we find that this formula is valid if and only if $\beta \geq \beta_0$, where β_0 is the real root of the polynomial

$$x^4 - 160x^3 + 3456x^2 + 55296x + 331776$$

whose approximate value is 42.04.

We prove $a(n + 1) \geq 43a(n) \geq 0$ for $n \geq 1$ by induction on n . The induction step follows from the formula above. As for the induction base, it suffices to check that $a(n + 1) \geq 43a(n) \geq 0$ for $n = 1, 2, 3, 4$, which is trivial.

The nonnegativity of $a(n)$ for $n \geq 1$ is hence established. Furthermore, $a(0) = 1 \geq 0$, so the proof is complete. \square

The proof for parts 2 and 3 of the theorem proceeds along the the same lines, of course with different recurrence equations, and consequently with different values for β_0 . Their approximate values for $r = 5$ and $r = 6$ are $\beta_0 \approx 138.9$ and $\beta_0 \approx 715.5$, respectively. We believe that for any specific value of r it is possible to obtain a similar proof, but the runtime requirements for the computations grow drastically and with currently available machines we were not able to go beyond $r = 6$ with reasonable effort. The runtime required for completing the cases $r = 4, 5, 6$ on a 2.4GHz Linux machine was, however, no more than a few minutes.

3. A RESULT OF SZEGÖ, AND A CONJECTURED VARIATION

Theorem 3. *Let $a(n, m, k)$ be such that*

$$\frac{1}{1 - (x + y + z) + \frac{3}{4}(xy + xz + yz)} = \sum_{n, m, k \geq 0} a(n, m, k)x^n y^m z^k.$$

Then $a(n, m, k) > 0$ for all $n, m \geq 0$ and (at least) $k = 0, 1, 2, \dots, 16$.

Proof. Because of symmetry it suffices to consider the case $n \geq m \geq k \geq 0$. For all the claimed values of k , we find a recurrence equation

$$p_{0,k}(n, m)a(n, m, k) + p_{1,k}(n, m)a(n + 1, m, k) + p_{2,k}(n, m)a(n + 2, m, k) = 0,$$

with $p_{i,k}(n, m)$ being polynomials in n and m for which the formula

$$\forall A_0, A_1, A_2 \in \mathbb{R} \forall n \geq m \geq k :$$

$$(A_1 \geq A_0 > 0 \wedge p_{0,k}(n, m)A_0 + p_{1,k}(n, m)A_1 + p_{2,k}(n, m)A_2 = 0) \implies A_2 \geq A_1$$

is true. This gives the induction step for proving $a(n + 1, m, k, l) \geq a(n, m, k, l) > 0$ for all $n \geq m \geq k \geq 0$. Checking the induction base is trivial. \square

We believe that the proof technique described above succeeds for every specific value of k ; again we were not able to verify this for $k > 16$ owing to the extensive runtime requirements of the algorithms.

Szegö [14] has shown by a rather complicated derivation that indeed $a(n, m, k) > 0$ for all $n, m, k \geq 0$. Later, Askey and Gasper [2] have given a different, but still complicated proof for the same fact. Both proofs rely on finding an integral representation for $a(n, m, k)$ and then applying arguments from the theory of special function for showing that the integrals are always positive.

Comparison to the simple recurrence equation obtained in Section 2 that asserts positivity of the coefficients $(1 - (x + y + z) + 4xyz)^{-1}$ leads naturally to the question whether a similar recurrence can be given for the coefficients of $(1 - (x + y + z) + \frac{3}{4}(xy + xz + yz))^{-1}$. No such recurrence has been published so far. By applying the procedure described in the proof of Theorem 1 to the present example, we have also not found such a recurrence. This, however, means that no such recurrence exists at all—a result that is perhaps not as easy to prove without computer assistance.

Proposition 1. *Let $a(n, m, k)$ be as in Theorem 3. Then there does not exist a recurrence equation*

$$\sum_{u, v, w \in \{0, 1\}} q_{u, v, w}(n, m, k)a(n + u, m + v, k + w) = 0 \quad (n, m, k \geq 0)$$

with linear polynomials $q_{u, v, w}(n, m, k)$ ($u, v, w \in \{0, 1\}$) such that

$$q_{1, 1, 1}(n, m, k) < 0 \text{ and } q_{u, v, w}(n, m, k) \geq 0 \text{ } ((u, v, w) \neq (1, 1, 1))$$

for all $n \geq m \geq k \geq 0$.

Despite some effort, we have also not been able to find a computer proof for general k by other means. We did find some simple recurrences for $a(n, m, k)$ but we did not succeed in constructing a positivity proof from any of them.

A variation of Theorem 3 arises as conjecture in the article of Askey and Gasper [2]: Are the coefficients in the series expansion of $(1 - (x + y + z + w) + \frac{2}{3}(xy + xz + xw + yz + yw + zw))^{-1}$ all positive? This conjecture is still open; the techniques of this paper are also insufficient for giving a complete proof. We can only offer additional evidence by supplying proofs for the situation where two indices are set to specific integers.

Theorem 4. *Let $a(n, m, k, l)$ be such that*

$$\frac{1}{1 - (x + y + z + w) + \frac{2}{3}(xy + xz + xw + yz + yw + zw)} = \sum_{n, m, k, l \geq 0} a(n, m, k, l)x^n y^m z^k w^l$$

Then $a(n, m, k, l) > 0$ for all $n, m \geq 0$ and (at least) all k, l with $0 \leq k + l \leq 14$.

Proof. Because of symmetry it suffices to consider the case $n \geq m \geq k \geq l \geq 0$. For all the claimed values of k and l , we find a recurrence equation

$$p_{0,k,l}(n, m)a(n, m, k, l) + p_{1,k,l}(n, m)a(n+1, m, k, l) + p_{2,k,l}(n, m)a(n+2, m, k, l) = 0$$

where $p_{i,k,l}(n, m)$ are some polynomials for which the formula

$$\forall A_0, A_1, A_2 \in \mathbb{R} \forall n \geq m \geq k :$$

$$(A_1 \geq A_0 > 0 \wedge p_{0,k,l}(n, m)A_0 + p_{1,k,l}(n, m)A_1 + p_{2,k,l}(n, m)A_2 = 0) \implies A_2 \geq A_1$$

is true. This gives the induction step for proving $a(n+1, m, k, l) \geq a(n, m, k, l) > 0$ for all $n \geq m \geq k \geq l \geq 0$. Checking the induction base is trivial. \square

Further evidence can be obtained by considering diagonals.

Theorem 5. *Let $a(n, m, k, l)$ be as in the previous theorem. Then $a(n, n+u, n+v, n+w) > 0$ for all $n \geq 0$ and (at least) all u, v, w with $0 \leq u, v, w \leq 12$.*

Proof. We abbreviate $a_{u,v,w}(n) := a(n, n+u, n+v, n+w)$. For all the claimed values of u, v, w , we find a recurrence equation

$$p_{0,u,v,w}(n)a_{u,v,w}(n) + p_{1,u,v,w}(n)a_{u,v,w}(n+1) + p_{2,u,v,w}(n)a_{u,v,w}(n+2) = 0$$

where $p_{i,u,v,w}(n)$ are some polynomials in n and m for which the formula

$$\forall A_0, A_1, A_2 \in \mathbb{R} \forall n \geq 3(u+v+w+2) :$$

$$(A_1 \geq \frac{64}{9}A_0 > 0 \wedge p_{0,u,v,w}(n)A_0 + p_{1,u,v,w}(n)A_1 + p_{2,u,v,w}(n)A_2 = 0) \implies A_2 \geq \frac{64}{9}A_1$$

is true. This gives the induction step for proving $a_{u,v,w}(n+1) \geq \frac{64}{9}a_{u,v,w}(n) > 0$. The proof is completed by checking $a_{u,v,w}(3(u+v+w+2)+1) \geq \frac{64}{9}a_{u,v,w}(3(u+v+w+2)) > 0$ as induction base, and $a_{u,v,w}(n) > 0$ for the points $0 \leq n < 3(u+v+w+2)$ not covered by the induction argument. \square

The mysterious constant $\frac{64}{9}$ that is needed in the induction step formula of the proof above was obtained in the same way as in the proof of Theorem 2. The lower bound $3(n+v+w+2)$ was found by experimenting and would probably have to be adjusted for values of u, v, w outside the range that we have considered.

4. TWO NEW CONJECTURES

We conclude this paper with two rational functions for which we conjecture that all their series coefficients are positive. Computational experiments have led us to these conjectures. As evidence in support of the conjectures, we provide partial proofs. We shall not state any opinion about the difficulty of proving the conjectures in full generality.

Theorem 6. *Let $a(n, m, k)$ be such that*

$$\frac{1}{1 - (x+y+z) + \frac{1}{4}(x^2+y^2+z^2)} = \sum_{n,m,k \geq 0} a(n, m, k)x^n y^m z^k.$$

Then $a(n, m, k) > 0$ for all $n, m \geq 0$ and (at least) $k = 0, 1, 2, \dots, 9$.

Proof. Because of symmetry it suffices to consider the case $n \geq m \geq k \geq 0$. For all the claimed values of k , we find a recurrence

$$p_{0,k}(n, m)a(n, m, k) + p_{1,k}(n, m)a(n+1, m, k) + p_{2,k}(n, m)a(n+2, m, k) = 0,$$

with $p_{i,k}(n, m)$ being polynomials in n and m for which the formula

$$\forall A_0, A_1, A_2 \in \mathbb{R} \forall n \geq m \geq 0 :$$

$$(A_1 \geq \frac{1}{2}A_0 > 0 \wedge p_{0,k}(n, m)A_0 + p_{1,k}(n, m)A_1 + p_{2,k}(n, m)A_2 = 0) \implies A_2 \geq \frac{1}{2}A_1.$$

is true. This is the induction step for proving $a(n+1, m, k) \geq \frac{1}{2}a(n, m, k) > 0$ for all $n \geq m \geq k \geq 0$. Checking the induction base is trivial. \square

Conjecture 1. Let $a(n, m, k)$ be as in Theorem 6. Then $a(n, m, k) > 0$ for all $n, m, k \geq 0$.

Computer experiments suggest furthermore that the conjecture becomes false if the constant $\frac{1}{4}$ in the denominator be replaced by $\frac{1}{4} + \varepsilon$ for any $\varepsilon > 0$.

Theorem 7. Let $a(n, m, k, l)$ be such that

$$\frac{1}{1 - (x + y + z + w) + \frac{64}{27}(xyz + xyw + xzw + yzw)} = \sum_{n, m, k, l \geq 0} a(n, m, k, l) x^n y^m z^k w^l.$$

Then $a(n, m, k, l) > 0$ for all $n, m \geq 0$ and (at least) all k, l with $0 \leq k + l \leq 12$.

Proof. For all the claimed values of k and l , we find that $a(n, m, k, l)$ is hypergeometric with respect to n , i.e., we find rational functions $r_{k,l}(n, m)$ such that

$$\forall n, m \geq 0 : a(n + 1, m, k, l) = r_{k,l}(n, m) a(n, m, k, l).$$

A CAD computation confirms that $r_{k,l}(n, m) > 0$ for all $n, m \geq k + l$, so positivity of the $a(n, m, k, l)$ follows from the positivity of $a(n, m, k, l)$ for $n, m \leq k + l$, which is easily verified. \square

Conjecture 2. Let $a(n, m, k, l)$ be as in Theorem 7. Then $a(n, m, k, l) > 0$ for all $n, m, k, l \geq 0$.

Also for this example, computer experiments suggest furthermore that the conjecture becomes false if the constant $\frac{64}{27}$ in the denominator be replaced by $\frac{64}{27} + \varepsilon$ for any $\varepsilon > 0$.

ACKNOWLEDGEMENT

The author is thankful to Doron Zeilberger for proposing the topic of this paper, and for his encouragement during its preparation.

REFERENCES

- [1] Horst Alzer, Stefan Gerhold, Manuel Kauers, and Alexandru Lupas. On Turán's inequality for Legendre polynomials. *Expositiones Mathematicae*, 2006. to appear.
- [2] Richard Askey and George Gasper. Certain rational functions whose power series have positive coefficients. *The American Mathematical Monthly*, 79(4):327–341, 1972.
- [3] Richard Askey and George Gasper. Convolution structures for Laguerre polynomials. *Journal d'Analyse Math.*, 31:46–68, 1977.
- [4] George E. Collins. Quantifier elimination for the elementary theory of real closed fields by cylindrical algebraic decomposition. *Lecture Notes in Computer Science*, 33:134–183, 1975.
- [5] Stefan Gerhold and Manuel Kauers. A procedure for proving special function inequalities involving a discrete parameter. In *Proceedings of ISSAC'05*, pages 156–162, 2005.
- [6] Stefan Gerhold and Manuel Kauers. A computer proof of Turán's inequality. *Journal of Inequalities in Pure and Applied Mathematics*, 7(2), 2006.
- [7] J. Gillis and J. Kleeman. A combinatorial proof of a positivity result. *Mathematical Proceedings of the Cambridge Philosophical Society*, 86:13–19, 1979.
- [8] J. Gillis, B. Reznick, and D. Zeilberger. On elementary methods in positivity theory. *SIAM Journal of Mathematical Analysis*, 14(2):396–398, 1983.
- [9] Manuel Kauers and Peter Paule. A computer proof of Moll's log-concavity conjecture. *Proceedings of the AMS*, 2006. to appear.
- [10] Tom Koornwinder. Positivity proofs for linearization and connection coefficients of orthogonal polynomials satisfying an addition formula. *Journal of the London Mathematical Society*, 18:101–114, 1978.
- [11] Peter Paule and Markus Schorn. A Mathematica version of Zeilberger's algorithm for proving binomial coefficient identities. *Journal of Symbolic Computation*, 20(5–6):673–698, 1995.
- [12] Bruno Salvy and Paul Zimmermann. Gfun: a Maple package for the manipulation of generating and holonomic functions in one variable. *ACM Transactions on Mathematical Software*, 20(2):163–177, 1994.
- [13] Adam Strzeboński. Solving systems of strict polynomial inequalities. *Journal of Symbolic Computation*, 29:471–480, 2000.
- [14] Gabor Szegő. Über gewisse Potenzreihen mit lauter positiven Koeffizienten. *Mathematische Zeitschriften*, 37(1):674–688, 1933.
- [15] Doron Zeilberger. The holonomic ansatz III: Towards automatic proofs of positivity results for coefficients of rational functions of several variables. in preparation.
- [16] Doron Zeilberger. The method of creative telescoping. *Journal of Symbolic Computation*, 11:195–204, 1991.

MANUEL KAUERS, RESEARCH INSTITUTE FOR SYMBOLIC COMPUTATION, J. KEPLER UNIVERSITY LINZ, AUSTRIA
E-mail address: `mkauers@risc.uni-linz.ac.at`