

Solving Difference Equations whose Coefficients are not Transcendental

Manuel Kauers¹

*Research Institute for Symbolic Computation
Johannes Kepler Universität
Altenberger Straße 69
A-4040 Linz, Austria, Europe*

Abstract

We consider a large class of sequences which are defined by systems of (possibly non-linear) difference equations. A procedure for recursively enumerating the algebraic dependencies of such sequences is presented. Also a procedure for solving linear difference equations with such sequences as coefficients is proposed. The methods are illustrated on some problems arising in the literature on special functions and combinatorial sequences.

Key words: Difference Equations, Algebraic Dependencies, Symbolic Summation

1 Introduction

A difference equation of order $r \in \mathbb{N}$ is an equation of the form

$$F(u(n), u(n+1), \dots, u(n+r), n) = 0 \quad (n \geq 1), \quad (1)$$

where $F: k^{r+2} \rightarrow k$ is an explicitly given function. Any sequence $u: \mathbb{N} \rightarrow k$ which satisfies (1) is called a solution of that equation. If F is linear in the first $r+1$ arguments, i.e., if the equation reads

$$a_0(n)u(n) + a_1(n)u(n+1) + \dots + a_r(n)u(n+r) = g(n) \quad (n \geq 1) \quad (2)$$

Email address: manuel.kauers@risc.uni-linz.ac.at (Manuel Kauers).

¹ Partially supported by the Austrian Science Foundation (FWF) grants F1305 and P19462-N18.

for some sequences $a_0, \dots, a_r, g: \mathbb{N} \rightarrow k$, then we call it a *linear difference equation*. Depending on the class of functions from which the a_i and g are chosen and on the class of functions in which solutions u are to be found, there are various known algorithms for solving linear difference equations. In the simplest case, the a_i and g are independent of n . In this case, a closed form solution in terms of exponentials can always be found by classical means [24]. Less trivial is the case where the a_i and g are rational functions in n . There are algorithms due to Abramov [2] and van Hoeij [36] which find all rational function solutions u to such equations. Petkovšek's algorithm [25] can find hypergeometric solutions u of the same type of equations. Hendriks and Singer [16] define the notion of liouvillean sequences and propose an algorithm for computing such solutions of linear difference equations with rational coefficients. Schneider [28] has got an algorithm for the case where the a_i may involve complicated expressions of nested sums and products. His algorithm finds solutions in terms of nested sums and products.

All these algorithms have in common that the coefficient sequences a_i can be written as rational functions of some basis sequences f_1, \dots, f_m which are algebraically independent (transcendental). In addition, these algorithms have the feature that they find *all* closed form solutions of a given equation in a finite number of steps. In the present paper, the focus is different. We aim at covering a very large class of linear difference equations, i.e., the class of functions, from which the functions a_i and g in (2) are chosen, is very rich. They are taken from a class of sequences introduced in 2003 by Zimmermann [38]. This class contains all the above-mentioned classes as subclasses, plus a lot of additional sequences, including in particular sequences that may obey non-trivial algebraic dependencies (Section 2). We will describe how to determine algebraic dependencies of such sequences, and how to find solutions of linear difference equations (2) whose coefficients a_0, \dots, a_r, g belong to this class. As we will argue, there is no reasonable hope that decision procedures for solving such problems exist. Instead, we will therefore propose procedures that recursively enumerate the set of all algebraic dependencies, respectively of all solutions to a given equation. Alternatively, these procedures may also be formulated as semi-decision procedures which, in a finite number of steps, find a solution if and only if there exists one, and otherwise run forever without producing any output.

Questions about difference equations need not be decidable. For example, the equivalence of systems of affine recurrence equations (SAREs) is undecidable [5]. There are several other problems for which it is not known whether they are decidable. For instance, whether a solution of a linear difference equation with constant coefficients has a root is not known to be decidable if the recurrence order exceeds five [12]. (The problem is decidable for small orders [15].) The situation in the differential case is similar. Here, we know various algorithms for finding closed form solutions of particular types of dif-

ferential equations [23,30,1,9,33,32,7]. For other types of differential equations, the existence of solutions is undecidable [4,11]. Also the existence of roots is undecidable already for a small class of continuous functions [27]. The zero equivalence problem is decidable for some classes of functions [29,34], but undecidable for some others [10]. For several problems in differential algebra, it is not known today whether or not they are decidable. For example it is not known whether the membership problem for differential ideals is decidable, while radical membership is known to be decidable [8]. It is also not known if there is an algorithm for deciding whether one prime differential ideal is contained in another [22]. To our knowledge, the corresponding questions in difference algebra are open as well.

The development of the algorithms described in the present paper was motivated by problems arising in the literature on special functions and combinatorial sequences, especially summation problems, which up to now could not be treated by symbolic computation. The overall goal is to devise methods for problems to which the classical algorithms are not applicable. Indeed, with the methods described in this paper, it is possible to solve some such problems. Example applications involving expression like F_{2^n} (the 2^n -th Fibonacci number) and other quantities, which are defined by nonlinear difference equations, are interspersed throughout the paper. In contrast to some of the algorithms mentioned in the beginning, our methods are conceptually simple and the mathematical background they are based on is rather moderate. Familiarity with basic facts of ideal theory and Gröbner basis techniques [6] is sufficient for reading this paper.

Not only are we interested in the theoretical aspects of solving difference equations. Rather, we claim that most of the proposed procedures are actually feasible on modern computer architectures. We are not able to justify this claim by any sort of theoretical statements about the complexity of our procedures. In fact, the worst case runtime of most algorithms is so poor that it is of little interest to actually determine bounds for their time and memory requirements. However, as it is often the case with algorithms related to commutative algebra, we observed that the computational cost for many examples is reasonably low, and certain identities arising in the literature could indeed be tackled by an implementation of the methods described in this article [19].

2 Admissible Sequences

In this paper, we consider sequences in a fixed computable field k of characteristic zero, i.e., functions $f: \mathbb{N} \rightarrow k$. The set of all sequences in k with pointwise addition and multiplication forms a commutative ring and is denoted by \mathcal{S} . Algorithms for sequences necessarily operate only on certain subsets of \mathcal{S} .

missible system

$$\left\{ f_1(n+1) = f_1(n) + 1, f_2(n+2) = \frac{2f_1(n) + 3}{f_1(n) + 2} x f_2(n+1) - \frac{f_1(n) + 1}{f_1(n) + 2} f_2(n) \right\}$$

with initial values $f_1(1) = 1$ (so that $f_1(n) = n$ for all n) and $f_2(1) = x$, $f_2(2) = \frac{1}{2}(3x^2 - 1)$ (so that $f_2(n) = P_n(x)$ for all n). Similarly, many other orthogonal polynomials, in fact all univariate holonomic sequences (also called P -finite sequences [37]) are admissible.

- (3) The sequence of Fibonacci numbers is admissible. Moreover, the sequence $n \mapsto F_{2^n}$ is admissible. To see this, recall that the Fibonacci numbers obey the addition theorems

$$\begin{aligned} F_{p+q} &= F_{p+1}F_q + F_pF_{q+1} - F_pF_q, \\ F_{p+q+1} &= F_pF_q + F_{p+1}F_{q+1}. \end{aligned}$$

Setting $p = q = 2^n$, we find

$$\begin{aligned} F_{2^{n+1}} &= F_{2^{n+2^n}} = F_{2^{n+1}}F_{2^n} + F_{2^n}F_{2^{n+1}} - F_{2^n}F_{2^n}, \\ F_{2^{n+1}+1} &= F_{2^{n+2^n}+1} = F_{2^n}F_{2^n} + F_{2^{n+1}}F_{2^{n+1}}, \end{aligned}$$

and consequently

$$\left\{ f_1(n+1) = 2f_2(n)f_1(n) - f_1(n)^2, f_2(n+1) = f_1(n)^2 + f_2(n)^2 \right\}$$

is a suitable admissible system for specifying the sequence $n \mapsto F_{2^n}$. By a similar construction, admissible systems for $n \mapsto 2^{F_n}$, $n \mapsto F_{F_n}$, and in fact for any sequence $n \mapsto f(g(n))$ where f and g satisfy homogeneous linear recurrences with constant coefficients and the coefficients in the recurrence of g are integral, can be obtained.

- (4) For some fixed $a_1, \dots, a_r \in k$, a sequence C satisfying the equation

$$\begin{aligned} C(n+r)C(n) &= a_1C(n+r-1)C(n+1) + a_2C(n+r-2)C(n+2) + \dots \\ &\quad \dots + a_rC(n+r-\lfloor r/2 \rfloor)C(n+\lfloor r/2 \rfloor) \end{aligned}$$

is an $(r-)$ Somos sequence [31, 13].

Somos sequences are admissible; a suitable admissible system is

$$\left\{ \begin{aligned} f_1(n+r) &= f_2(n) \left(a_1f_1(n+r-1)f_1(n+1) + \dots \right. \\ &\quad \left. \dots + a_rf_1(n+r-\lfloor r/2 \rfloor)f_1(n+\lfloor r/2 \rfloor) \right), \\ f_2(n) &= 1/f_1(n) \end{aligned} \right\}$$

Observe that the sequence f_2 was introduced to fulfill the requirement of Def. 1 that numerators or denominators of the rational functions on the right hand side be constant.

It should be remarked at this point that for some admissible systems, not every choice of initial values yields well-defined sequences. This is because denominators might become zero for some points. For instance, the admissible system $\{f_1(n+1) = f_1(n) + 1, f_2(n) = 1/f_1(n)\}$ defines two admissible sequences f_1, f_2 once the initial value $f_1(1)$ is chosen. If a negative integer is chosen for $f_1(1)$, then $f_2(n)$ is undefined at $n = -f_1(1)$. For admissible systems and initial values which are supplied as input of our algorithms, we will always assume that this situation does not occur, i.e., that the input sequences are well-defined.

New admissible sequences can be composed out of known ones by using the following closure properties of the class of admissible sequences.

Theorem 3 [18, Thms. 3.5, 3.7] *Let f and g be admissible sequences, $a \in \mathbb{N}$ and $\alpha \in k$. Then*

- (1) αf , $f + g$ and $f \cdot g$ and, if $g(n) \neq 0$ for all n , f/g are admissible,
- (2) $n \mapsto \sum_{i=1}^n f(i)$, $n \mapsto \prod_{i=1}^n f(i)$, and, if $f(n) \neq 0 \neq g(n)$ for all n ,

$$n \mapsto f(1) + \frac{g(2)}{f(2) + \frac{g(3)}{\dots + \frac{g(n)}{f(n)}}$$

are admissible,

- (3) $n \mapsto f(n+a)$, $n \mapsto f(an)$, $n \mapsto f(\lfloor n/a \rfloor)$ are admissible,

and admissible systems for these sequences can be effectively computed from admissible systems for f and g .

Using this theorem, it is possible to automatically transform an expression involving sums and products into a corresponding defining admissible system. Rather than giving a formal proof, we illustrate the theorem with an example.

Example 4 *The expression*

$$F_n^2 \sum_{k=1}^{n-1} \frac{1}{F_k^2 F_{k+1}^2} \prod_{i=2}^k \frac{F_{i-1}}{2F_i - F_{i-1}}$$

constitutes an admissible sequence in \mathbb{Q} . $(0, 0, 1, 5, \frac{34}{3}, \frac{63}{2}, \dots)$ A suitable admissible system can be constructed by first considering the innermost subex-

pressions and then building up the whole expression step by step:

$$\left\{ \begin{array}{ll} f_1(n+2) = f_1(n+1) + f_1(n) & (f_1(n) \sim F_n) \\ f_2(n+1) = 1/(2f_1(n+1) - f_1(n)) & (f_2(n) \sim 1/(2F_n - F_{n-1})) \\ f_3(n+1) = f_3(n)f_1(n)f_2(n) & (f_3(n) \sim \Pi) \\ f_4(n+1) = 1/(f_1(n)^2 f_1(n+1)^2) & (f_4(n) \sim 1/F_n^2 F_{n-1}^2) \\ f_5(n+1) = f_5(n) + f_4(n+1)f_3(n) & (f_5(n) \sim \Sigma) \\ f_6(n+1) = f_1(n+1)^2 f_5(n) \end{array} \right\} \quad (f_6(n) \sim \text{whole expr.})$$

It is easily seen that the class of admissible sequences properly includes many of the classes known for the algorithms mentioned in the introduction, such as the (univariate) holonomic sequences [37] and Karr's $\Pi\Sigma$ sequences [17], for instance. In addition, sequences like $n \mapsto F_{F_n}$, which are admissible, do not belong to any class of sequences that can be handled by a known algorithm. Yet there are — of course — still sequences which are not covered. For instance, it can be shown [18, Section 4.3] that the sequences $n \mapsto (-1)^{\lfloor \log n \rfloor}$ and $n \mapsto 2^{n!}$ cannot be defined via an admissible system.

3 Reduction to Polynomial Algebra

Let f_1, \dots, f_m be admissible sequences, and consider the ring homomorphism $\phi: k[x_1, \dots, x_m] \rightarrow \mathcal{S}$ that maps x_i to f_i and each $c \in k$ to the constant sequence (c, c, c, \dots) . The homomorphism theorem asserts that the factor ring $k[x_1, \dots, x_m]/\ker \phi$ is isomorphic to $\text{im } \phi$, which is the smallest subring of \mathcal{S} containing f_1, \dots, f_m and all constant sequences. As $\ker \phi$ is just a polynomial ideal, the computational treatment of the ring $k[x_1, \dots, x_m]/\ker \phi$ is well understood. The theory of Gröbner bases [6] provides an algebraic framework for solving problems in such domains. By the isomorphism, there is a one-to-one correspondence between $k[x_1, \dots, x_m]/\ker \phi$ and $\text{im } \phi$, so that results obtained in the former ring (by Gröbner basis or other means) can be directly translated into results in the latter ring, which is in our interest.

There is only a slight obstacle here: in order to perform computations in the ring $k[x_1, \dots, x_m]/\ker \phi$, we need to know some rather explicit information about the ideal $\ker \phi$, for instance a list of ideal generators $p_1, \dots, p_s \in k[x_1, \dots, x_m]$ such that $\ker \phi = \langle p_1, \dots, p_s \rangle$. Ideally, we would like to be able to compute such generators p_1, \dots, p_s from a given admissible system and initial values for the f_1, \dots, f_m . No algorithms are known for this problem. However, in an earlier paper [20] we have shown that the membership problem for $\ker \phi$ (given $p \in k[x_1, \dots, x_m]$, decide $p \in \ker \phi$) is decidable. The remarkable aspect of this algorithm is that generators of $\ker \phi$ need not be known; only an

admissible system for the f_1, \dots, f_m and initial values are required as input.

Note that $p \in \ker \phi$ just means that $F := p(f_1, \dots, f_m)$ is the zero sequence. Regardless of whether this is the case, F is for sure an admissible sequence, because the f_i are admissible (Thm. 3). Our decision procedure [20] is hence an algorithm for deciding zero equivalence of admissible sequences. It works by constructing an induction proof for the sequence F to be zero. In the first phase, it computes a number $N \in \mathbb{N}$ with the property that

$$\forall n \in \mathbb{N} : F(n) = F(n+1) = \dots = F(n+N-1) = 0 \Rightarrow F(n+N) = 0.$$

This N provides the induction step. In the second phase, the algorithm evaluates $F(1), F(2), \dots, F(N)$ and determines whether they are all zero. (It is assumed that the ground field k is such that zero equivalence in the ground field can be decided.) This either supplies the base of the induction and thus a decision that indeed $F \equiv 0$, or it leads to an index $n \in \{1, \dots, N\}$ with $F(n) \neq 0$ and thus a decision that $F \not\equiv 0$. The algorithm reveals the following theorem.

Theorem 5 [20,18] *There exists an algorithm which for a given admissible system S and $p \in k[x_1, \dots, x_m]$ computes a number $N \in \mathbb{N}$ such that for all solutions f_1, \dots, f_m of S :*

$$\forall n \in \mathbb{N} : F(n) = 0 \iff F(1) = F(2) = \dots = F(N) = 0,$$

where $F = p(f_1, \dots, f_m)$. \square

It must be remarked that the number N asserted by the above theorem only depends on the admissible system S , but not on the initial values of the f_1, \dots, f_m . For further details about this algorithm, we refer to [20,18].

4 Recursive Enumeration of a Basis for the Kernel

Our goal is to find elements of the ideal $\ker \phi$. There is little hope that an algorithm could be found which computes a basis of $\ker \phi$ from an admissible system and initial values for the f_1, \dots, f_m . This is because of the following reduction.

Theorem 6 *If there exists an algorithm which for given admissible sequences f_1, \dots, f_m computes a basis of $\ker \phi$, then there exists an algorithm which for a given admissible sequence f decides whether there exists an index n with $f(n) = 0$, and, if yes, delivers the smallest such n .*

Proof. Let f be an admissible sequence, say f is defined via an admissible system for f_1, \dots, f_m and $f = f_i$. If we define $f_0(n) := n$ and $f_{m+1}(n) =$

$\prod_{j=1}^n f_i(j)$, then f_0 and f_{m+1} are admissible, too, by Theorem 3.

Let $\phi: k[x_0, \dots, x_{m+1}] \rightarrow \mathcal{S}$ be defined via $\phi(x_i) = f_i$ ($i = 0, \dots, m+1$). Then $f(n_0) = 0$ for some $n_0 \in \mathbb{N}$ if and only if

$$x_{m+1}(x_0 - 1)(x_0 - 2) \cdots (x_0 - n_0) \in \ker \phi.$$

Suppose now that a basis $\ker \phi = \langle p_1, \dots, p_m \rangle$ is known. Then we can compute a Gröbner basis G for this ideal with respect to the lexicographic order $x_{m+1} > x_m > \cdots > x_1 > x_0$. This G contains the polynomial $x_{m+1}(x_0 - 1)(x_0 - 2) \cdots (x_0 - n_0)$ indicating the smallest root n_0 of f — or no polynomial of this form if f does not have any roots. \square

According to this theorem, the computation of a basis for $\ker \phi$ is at least as difficult as finding a root of an admissible sequence. Unfortunately, finding roots of sequences is a very difficult problem. Already for the class of sequences satisfying homogeneous linear recurrence equations with constant coefficients (also called C-finite sequences [37]), it is an open problem whether the question about the existence of a root is decidable [12, Sec. 2.3]. As this class is only a very small subclass of the class of admissible sequences, it does not seem reasonable to look for an algorithm that would be capable of solving this problem for the much larger class. We want to describe instead a procedure by which a basis of the kernel $\ker \phi$ can be recursively enumerated.

4.1 Linear Dependencies of Admissible Sequences

As a subalgorithm for the enumeration procedure, we need an algorithm for computing linear dependencies of admissible sequences. Given a finite set of polynomials, $P = \{p_1, \dots, p_l\} \subseteq k[x_1, \dots, x_m]$, this algorithm computes a basis of the vector space

$$V_P := (p_1k + p_2k + \cdots + p_lk) \cap \ker \phi \subseteq k[x_1, \dots, x_m].$$

The algorithm proceeds by making an undetermined ansatz, obtaining candidates for the desired relations by comparing them with the initial values and solving a linear system. The candidates can be validated by the algorithm of Theorem 5. If necessary, the ansatz is refined more and more, until all candidates actually belong to the kernel.

Algorithm 7 Input: Admissible sequences $f_1, \dots, f_m: \mathbb{N} \rightarrow k$, a set $P = \{p_1, \dots, p_l\} \subseteq k[x_1, \dots, x_m]$

Output: A basis of the vector space V_P .

- 1 Define $g_i := p_i(f_1, \dots, f_m)$ ($i = 1, \dots, m$)

2 $N = l$;
 3 **repeat**
 4 $N = N + 1$
 5 Compute a basis $B \subseteq k^l$ for the solution space of the linear system

$$\begin{pmatrix} g_1(1) & \cdots & g_l(1) \\ \vdots & \ddots & \vdots \\ g_1(N) & \cdots & g_l(N) \end{pmatrix} \begin{pmatrix} c_1 \\ \vdots \\ c_l \end{pmatrix} = 0$$

6 **until** $b_1g_1(n) + b_2g_2(n) + \cdots + b_lg_l(n) = 0$ ($n \in \mathbb{N}$) for all $(b_1, \dots, b_l) \in B$
 7 **return** $\{ b_1p_1 + b_2p_2 + \cdots + b_l p_l : (b_1, \dots, b_l) \in B \}$

Observe that the g_i are admissible sequences by Theorem 3(1). Therefore, the condition in line 6 can be decided according to Theorem 5.

Theorem 8 *Algorithm 7 is correct, i.e., if B is the set of polynomials returned by the algorithm, then B is a basis of the vector space V_P .*

Proof. Obviously, each element of B belongs to V_P by line 6, and by construction the elements of B are linearly independent. It only remains to show that every $c_1p_1 + \cdots + c_l p_l \in V_P$ is a linear combination of the vectors in B . For every vector $c_1p_1 + \cdots + c_l p_l \in V_P$ the identity

$$c_1g_1(n) + \cdots + c_lg_l(n) = 0$$

holds for all $n \geq 1$, by definition of V_P . In particular it holds for $n = 1, \dots, N$, and hence (c_1, \dots, c_l) belongs to the solution space of the linear systems in line 5. \square

Theorem 9 *Algorithm 7 terminates, i.e., in the notation of the algorithm, for sufficiently large N , all elements of B will give rise to kernel elements.*

Proof. Let S be an admissible system defining f_1, \dots, f_m , and consider the admissible system

$$S' := S \cup \{ f_{m+1}(n+1) = f_{m+1}(n), \dots, f_{m+l}(n+1) = f_{m+l}(n) \}$$

for f_1, \dots, f_{m+l} . Let $P := f_{m+1}p_1(f_1, \dots, f_m) + \cdots + f_{m+l}p_l(f_1, \dots, f_m)$. By Theorem 5, there exists a number $N \in \mathbb{N}$ such that $P \equiv 0$ if and only if $P(1) = P(2) = \cdots = P(N) = 0$. This N bounds the number of iterations in the loop in Algorithm 7. \square

4.2 Algebraic Dependencies of Admissible Sequences

It is a simple matter to extend Algorithm 7 to the desired enumeration procedure: just apply the algorithm in turn to find all linear dependencies of all the polynomials with total degree $d = 1, 2, 3, \dots$. The union of the outputs for all $d \in \mathbb{N}$ is obviously a k vector space basis for $\ker \phi$. Unless $\ker \phi = \{0\}$, this basis will be infinite. The vector space basis is also an ideal basis, but a rather redundant one. In order to obtain an irredundant ideal basis, we should restrict the set P in the input of Algorithm 7 in such a way that solutions of the linear system are not already consequences of the dependencies accumulated for degrees smaller than d . This can be done as follows.

Procedure 10 *Input:* Admissible sequences $f_1, \dots, f_m: \mathbb{N} \rightarrow k$

Output: An ideal basis of $\ker \phi$

- 1 $G = \emptyset; d = 0$
- 2 **repeat**
- 3 Let P be a vector space basis of

$$\{p \in k[x_1, \dots, x_m] : \deg p \leq d\}$$
- 4 Delete from P all elements p with $\text{LT}(g) \mid \text{LT}(p)$ for some $g \in G$
- 5 Apply Algorithm 7 to f_1, \dots, f_m and P , obtaining B
- 6 Output the elements of B
- 7 $G = \text{GröbnerBasis}(G \cup B)$
- 8 $d = d + 1$

To be specific, assume that $k[x_1, \dots, x_m]$ is equipped with a total degree term order. Any other admissible term order [6, Def. 4.59] could be taken instead. By $\text{LT}(p)$ we mean the leading term of a polynomial $p \in k[x_1, \dots, x_m]$ with respect to that order.

Theorem 11 *Procedure 10 is correct, i.e., the polynomials it outputs generate $\ker \phi$ as an ideal.*

Proof. Without line 4, the Theorem would be evident. We have to show that no relations are lost in line 4. In other words, if \mathfrak{a} denotes the ideal generated by the output of the procedure, we have to show that $\ker \phi \subseteq \mathfrak{a}$.

Suppose for the contrary that there exists $p \in \ker \phi \setminus \mathfrak{a}$. Then, because Algorithm 7 is correct, $\text{LT}(p)$ must be a multiple of $\text{LT}(a)$ for some $a \in \mathfrak{a}$. The leading term of $p' := p - \frac{\text{LM}(p)}{\text{LM}(a)}a$ (with LM being the leading monomial) is smaller than that of p , and p' also belongs to $\ker \phi \setminus \mathfrak{a}$, because $p \notin \mathfrak{a}$ and $a \in \mathfrak{a} \subseteq \ker \phi$. Repeating the argument, we find $p'' \in \ker \phi \setminus \mathfrak{a}$ with a leading

term smaller than that of p' , and so on. This leads to an infinite descending chain of terms $\text{LT}(p), \text{LT}(p'), \text{LT}(p''), \dots$, which by the admissibility of the term order cannot exist. \square

Example 12 Consider the sequences f_1, f_2, f_3 defined by

$$f_1(n) = F_{2^{n+a}}, \quad f_2(n) = F_{2^{n+a+1}}, \quad f_3(n) = \sum_{k=0}^n \frac{1}{F_{2^{a+k}}}.$$

Applying Procedure 10, we find no algebraic dependencies of total degrees 0, 1, 2 between these sequences, i.e., there does not exist a polynomial $p \in \mathbb{Q}[x, y, z]$ of total degree at most 2 with $p(f_1, f_2, f_3) \equiv 0$. For degree 3, the procedure delivers the relations

$$\begin{aligned} 0 &= F_{2^{n+a}} F_{2^{2n+a+1}}^2 - F_{2^{2n+a+1}}^3 + F_{2^{2n+a+1}}^2 + F_{2^{2n+a}}^2 F_{2^{n+a+1}} \\ &\quad - F_{2^{n+a}} F_{2^{n+a+1}} + F_{2^{n+a+1}} - F_{2^{2n+a}} - 1, \\ 0 &= F_{2^{n+a}}^3 - 2F_{2^{n+a+1}}^2 F_{2^{n+a}} + F_{2^{n+a}} + F_{2^{2n+a+1}}^3 - F_{2^{n+a+1}} \end{aligned}$$

therefore $\langle xy^2 - y^3 + y^2 + x^2y - xy + y - x^2 - 1, x^3 - 2y^2x + x + y^3 - y \rangle \subseteq \ker \phi$, with $\phi: \mathbb{Q}[x, y, z] \rightarrow \mathcal{S}$ such that $\phi(x) = f_1, \phi(y) = f_2, \phi(z) = f_3$. For total degree 4, 5, 6, \dots , 10, there are no further relations. Observe that the ideal generated by the relations we found may be written as $\langle x-1, y-1 \rangle \cap \langle 1+x^2+xy-y^2 \rangle$, so for each $n \in \mathbb{N}$ we have $f_1(n) = f_2(n) = 1$ or $f_1(n)^2 + f_1(n)f_2(n) - f_2(n)^2 = -1$.

Procedure 10 does not terminate. However, after a finite number of steps, it will have output a complete basis of $\ker \phi$. This follows from Hilbert's basis theorem: If \mathfrak{a}_d denotes the ideal generated by the output of the first d iterations ($d = 0, 1, 2, \dots$), then

$$\mathfrak{a}_0 \subseteq \mathfrak{a}_1 \subseteq \mathfrak{a}_2 \subseteq \dots$$

is an infinite ascending chain of polynomial ideals, so there must be some index d_0 such that $\mathfrak{a}_d = \mathfrak{a}_{d_0}$ for all $d \geq d_0$. As the procedure in each iteration $d > 0$ only outputs elements of $\mathfrak{a}_d \setminus \mathfrak{a}_{d-1}$ (this is easy to see), it follows that for $d \geq d_0$, no further output will happen. What is missing for a full algorithm is a way to compute a suitable upper bound for the value d_0 .

A finite algorithm for computing a basis can be obtained by restricting the attention to smaller classes of sequences. For instance, Karr's summation algorithm [17] includes as a subroutine an algorithm for computing the algebraic dependencies of sequences which can be expressed in terms of nested sums and products. Also the algebraic dependencies of sequences which satisfy homogeneous linear difference equations with constant coefficients can be effectively computed [21,18]. It would be desirable to further investigate for which classes of sequences the algebraic dependencies are computable.

5 Applications

The procedures introduced in the previous section enable us to solve certain problems appearing in the literature on special functions and combinatorial sequences automatically. In this section, we want to illustrate some applications with concrete examples.

Example 13 Consider the 4-Somos sequence $C(n)$ defined by

$$\begin{aligned} C(n+2) &= (C(n-1)C(n+1) + C(n)^2)/C(n-2) \quad (n \in \mathbb{Z}), \\ C(-2) &= C(1) = C(0) = C(1) = 1. \end{aligned}$$

It is of interest to know whether $C(n)$ also satisfies an r -Somos recurrence for some $r \neq 4$ (cf. [35]).

For any given r , say $r = 8$, this question can be answered using Algorithm 7: compute a vector space basis $\{b_1, \dots, b_s\}$ of

$$\ker \phi \cap \{p \in \mathbb{Q}[x_{-r}, x_{-r+1}, \dots, x_r] : \deg p \leq 2\},$$

where $\phi: \mathbb{Q}[x_{-r}, \dots, x_r] \rightarrow \mathcal{S}$ maps x_i to the sequence $n \mapsto C(n+i)$. For $r = 8$, this basis is lengthy and not reproduced here.

We have to find out whether the b_i can be combined to a relation of the desired form. One way to do so is to make an ansatz

$$a_4 C(n-4)C(n+4) + a_3 C(n-3)C(n+3) + \dots + a_0 C(n)^2 = 0$$

for the coefficients a_i , compute the normal form of the polynomial $a_4 x_{-4} x_4 + a_3 x_{-3} x_3 + a_2 x_{-2} x_2 + a_1 x_{-1} x_1 + a_0 x_0^2$ with respect to the ideal $\langle b_1, \dots, b_s \rangle$ (and some term order), equate the coefficients of that normal form to zero and solve the resulting linear system for the a_i . The solutions of this system are precisely the desired values for the coefficients.

In this way, we have found the relations

$$\begin{aligned} C(n+2)C(n-2) &= C(n+1)C(n-1) + C(n)^2 \\ C(n+3)C(n-3) &= C(n+1)C(n-1) + 5C(n)^2 \\ C(n+4)C(n-4) &= 25C(n+1)C(n-1) - 4C(n)^2, \end{aligned}$$

and by an analogous ansatz for odd r the relations

$$\begin{aligned} C(n+3)C(n-2) &= 5C(n+1)C(n) - C(n+2)C(n-1) \\ C(n+4)C(n-3) &= C(n+1)C(n-1) + 5C(n)^2. \end{aligned}$$

The first three relations were also given by van der Poorten [35], the last one is new. By Theorem 8, we can be sure that every other r -Somos recurrence of C for $r \leq 8$ is a linear combinations of those given above.

Example 14 Certain nonlinear difference equations can be solved using Procedure 10. For instance, Rabinowitz [26] has asked for a solution of

$$u(n+1) = \frac{3u(n) + 1}{5u(n) + 3} \quad (n \geq 1), \quad u(1) = 1$$

in terms of Fibonacci numbers. If there exists a rational function $r = p/q$ with $p, q \in \mathbb{Q}[x, y]$ such that $u(n) = r(F_n, F_{n+1})$ for all $n \geq 1$, then $q(x, y)z - p(x, y) \in \ker \phi$, where $\phi: \mathbb{Q}[x, y, z] \rightarrow \mathcal{S}$ maps x to the Fibonacci sequence $n \mapsto F_n$, y to the shifted Fibonacci sequence $n \mapsto F_{n+1}$, and z to u .

In order to find a solution, we apply Procedure 10. After each iteration, we compute a lexicographic Gröbner basis G of \mathfrak{a} with respect to $z > y > x$ and check whether G contains a polynomial linear in z . Each such polynomial supplies a solution, and if no such polynomial appears in G then no such polynomial is contained in \mathfrak{a} , and we increase the degree.

This procedure will eventually reveal any solution of the difference equation in terms of Fibonacci numbers—if such a solution exists at all. Otherwise, the procedure will run forever. In the present situation, we find

$$u(n) = -\frac{2F_n^2 - 2F_nF_{n+1} + F_{n+1}^2}{4F_n^2 - 6F_nF_{n+1} + F_{n+1}^2} \quad (n \geq 1).$$

By leaving the initial value $u(1)$ symbolic during the computation, the more general solution

$$u(n) = \frac{(3 - 7u(1))F_n^2 - 4(2u(1) - 1)F_nF_{n+1} + (1 - 3u(1))F_{n+1}^2}{(15u(1) - 7)F_n^2 + 4(2 - 5u(1))F_nF_{n+1} + (5u(1) - 3)F_{n+1}^2}$$

can be found.

Example 15 Linear difference equation can be treated as explained for nonlinear equations in the previous example. This includes indefinite summation as a special case. The important aspect here is that any admissible sequence may occur in the summand. Identities like

$$\sum_{k=0}^n \frac{1}{F_{2^k}} = 4 - \frac{F_{2^{n+1}}}{F_{2^n}}, \quad \sum_{k=0}^n \frac{1}{F_{3 \cdot 2^k}} = \frac{9}{4} - \frac{F_{3 \cdot 2^{n+1}}}{F_{3 \cdot 2^n}} \quad (n \geq 1)$$

[14, Ex. 6.61] can thus be found automatically. In contrast, no closed form for $\sum_{k=0}^n 1/F_{2^{k+a}}$ in terms of $F_{2^{n+a}}$ and $F_{2^{n+a+1}}$ is found (cf. Example 12).

6 Linear Difference Equations

Linear difference equations deserve special attention because of their importance in practice. Although we could find solutions of linear difference equations by means of Procedure 10 just as explained before for nonlinear difference equations, we would like to describe an alternative method for this special case. Let us consider an equation of the form

$$a_r(n)u(n+r) + a_{r-1}(n)u(n+r-1) + \cdots + a_0(n)u(n) = g(n), \quad (3)$$

where a_0, \dots, a_r, g are known admissible sequences and u is unknown. If a_r has finitely many roots only, then u is determined uniquely by finitely many initial values, and in particular every solution is admissible. Otherwise, if $a_r(n)$ has infinitely many solutions, then the equation has a continuum of solutions.

What interests us here is not the general solution of equation 3, but solutions of a prescribed form. We will assume that admissible sequences f_1, \dots, f_m are given and that solutions of (3) are to be computed which have the form $p(f_1, \dots, f_m)$ for some polynomial p .

6.1 The Homogeneous Equation

Let us first consider the case of a homogeneous equation, $g(n) = 0$. In order to find polynomials p such that $u := p(f_1, \dots, f_m)$ satisfies (3), we first use Procedure 10 to compute generators of the ideal $\ker \phi$ where

$$\phi: \underbrace{k[x_{1,0}, \dots, x_{m,0}, \dots, x_{1,r}, \dots, x_{m,r}, y_0, \dots, y_r]}_{=:R} \rightarrow \mathcal{S}$$

maps $x_{i,j}$ to $n \mapsto f_i(n+j)$, y_i to $n \mapsto a_i(n)$, and constants to constants. Assuming that $\ker \phi$ is known, we then compute a basis of the syzygy module

$$S := \text{Syz}(y_0, \dots, y_r) = \{ (p_0, \dots, p_r) : p_0 y_0 + \cdots + p_r y_r = 0 \} \subseteq (R/\ker \phi)^{r+1}.$$

It is well known how to compute the syzygy module over a polynomial ring [6, Sec. 6.1], and it is straightforward to generalize this algorithm to the case of a factor ring $k[X]/\mathfrak{a}$ [18, Thm. 2.9]. (For simplicity of notation, we will not distinguish polynomials and their residue classes modulo $\ker \phi$.) Now observe that for every polynomial $p \in k[x_1, \dots, x_m]$ we have

$$\begin{aligned} p(f_1, \dots, f_m) \text{ solves (3)} \\ \iff \\ (p(x_{1,0}, \dots, x_{m,0}), \dots, p(x_{1,r}, \dots, x_{m,r})) \in S \end{aligned} \quad (4)$$

Hence, we can find polynomials p with the desired property as follows. If we make an ansatz $p = \sum_{i_1, \dots, i_m} a_{i_1, \dots, i_m} x_1^{i_1} \cdots x_m^{i_m}$ for the solution polynomial and compute the normal form of the vector

$$\left(p(x_{1,0}, \dots, x_{m,0}), \dots, p(x_{1,r}, \dots, x_{m,r}) \right)$$

for this general p with respect to a Gröbner basis of S , then we will end up with some vector (q_0, \dots, q_r) where each q_i is a polynomial whose coefficients are linear combinations of the as yet undetermined a_{i_1, \dots, i_m} . The ansatz polynomial p represents a solution precisely for those values a_{i_1, \dots, i_m} that make all q_i vanish, because of (4) and the fact that normal forms are zero precisely for vectors which belong to the module. Comparing the coefficients of the q_i to zero gives rise to a linear system over k for the coefficients a_{i_1, \dots, i_m} which can be solved.

Algorithm 16 *Input:* A Gröbner basis G of a module $S \subseteq (R/\mathfrak{a})^s$, a set $P = \{p_1, \dots, p_l\} \subseteq (R/\mathfrak{a})^s$

Output: A basis for the vector space of all linear combinations p of p_1, \dots, p_l with $p \in S$

- 1 Make an ansatz $p = a_1 p_1 + \cdots + a_l p_l$
- 2 Compute the normal form (q_1, \dots, q_s) of p w.r.t. G
- 3 Let $c_i(a_1, \dots, a_l)$ ($i \in I$) be the coefficients of q_0, \dots, q_s
- 4 Compute a basis B of the space $\{ (a_1, \dots, a_l) : c_i(a_1, \dots, a_l) = 0 \ (i \in I) \}$
- 5 **return** $\{ a_1 p_1 + \cdots + a_l p_l : (a_1, \dots, a_l) \in B \}$

Termination of this algorithm is obvious, and its correctness follows from the discussion above. Applying the algorithm in turn to bigger and bigger ansatz polynomials, we obtain a procedure that recursively enumerates a basis for the solution space of (3). We cannot hope for a termination criterion (such as, e.g., a degree bound) here either, because the solution space may have infinite dimension. As in Procedure 10, we can discard leading terms to avoid redundant solutions and to keep the linear systems small. Cancellation of leading terms also ensures the output solutions are linearly independent.

Procedure 17 *Input:* Admissible sequences a_0, \dots, a_r and f_1, \dots, f_m and a basis of $\ker \phi$ with $\phi: R \rightarrow \mathcal{S}$ as defined above.

Output: A basis of the vector space of all solutions u of

$$a_0(n)u(n) + \cdots + a_r(n)u(n+r) = 0$$

which depend polynomially on f_1, \dots, f_m .

- 1 $B = \emptyset; d = 0$
- 2 Let G be a Gröbner basis of $\ker \phi$

3 Let S be a Gröbner basis of $\text{Syz}(y_0, \dots, y_r) \subseteq (R/\ker \phi)^{r+1}$
4 **repeat**
5 Let P be a vector space basis of $\{p \in k[x_1, \dots, x_m] : \deg p \leq d\}$
6 Delete from P all elements p with $\text{LT}(g) \mid \text{LT}(p)$ for some $g \in G$
7 Delete from P all elements p with $\text{LT}(b) = \text{LT}(p)$ for some $b \in B$
8 $P := \{ (p(x_{1,0}, \dots, x_{m,0}), \dots, p(x_{1,r}, \dots, x_{m,r})) : p \in P \}$
9 Apply Algorithm 16 to P and S , obtaining B_0
10 $B_0 := \{ q_0 : (q_0, \dots, q_r) \in B_0 \}$
11 Output the elements of B_0
12 $B := B \cup B_0; d = d + 1$

Theorem 18 *Procedure 17 is correct, i.e., its output constitutes a vector space basis of the k vector space of all solutions u of (3) which can be written polynomially in f_1, \dots, f_m .*

Proof. First of all, it is clear that every output polynomial really gives rise to a solution. We have to show that (a) no solutions are overlooked due to lines 6 and 7 and (b) the output solutions are linearly independent over k .

(a) Let p be a polynomial that corresponds to a solution u of the difference equation. The polynomial p is equivalent modulo $\ker \phi$ to a polynomial p' that does not contain terms which are multiples of leading terms in G . This polynomial p' corresponds to the same solution u so it suffices to take the terms into account that may possibly occur in p' .

Secondly, if a solution polynomial p involves a term τ which appears as a leading term of some solution $b \in B$ which was found before, then $p' := p - \alpha b$ for a suitable constant $\alpha \in k$ is another solution which does not involve τ . Restricting the ansatz such that only p' is found is just fine, because p is a linear combination of p' and b .

(b) Induction to d . For $d = 0$, $B = \emptyset$ is linearly independent. Now suppose that B is linearly independent at iteration d and assume $b_1, \dots, b_v \in B$, $c_1, \dots, c_w \in B_0$, and $\beta_1, \dots, \beta_v, \gamma_1, \dots, \gamma_w \in k$ are such that

$$\beta_1 b_1 + \dots + \beta_v b_v + \gamma_1 c_1 + \dots + \gamma_w c_w = 0. \quad (5)$$

If $\beta_1 b_1 + \dots + \beta_v b_v$ is not the zero polynomial, then we must have

$$\text{LT}(\beta_1 b_1 + \dots + \beta_v b_v) = \text{LT}(\gamma_1 c_1 + \dots + \gamma_w c_w),$$

which is excluded by line 7. It follows that $\beta_1 b_1 + \dots + \beta_v b_v = 0$, and hence by (5) also $\gamma_1 c_1 + \dots + \gamma_w c_w = 0$. Now using the linear independence of B and B_0 , respectively, we obtain $\beta_1 = \dots = \beta_v = \gamma_1 = \dots = \gamma_w = 0$, as desired. \square

Procedure 17 requires knowledge of $\ker \phi$ as input, but we only know a recursive enumeration procedure for computing $\ker \phi$ (Procedure 10). If in practice that procedure is aborted after a while, then it is not clear whether the ideal \mathfrak{a} generated by the output produced before abortion already generates the whole ideal $\ker \phi$. It is therefore interesting to know to what extent Procedure 17 remains correct if it is applied to some ideal $\mathfrak{a} \subsetneq \ker \phi$ in place of $\ker \phi$. It is quite easy to see that its output will still be correct and complete, but the output might be redundant. The sequence of polynomials it produces will also continue to be linearly independent, but this need no longer be true for the sequence of solutions u that these polynomials represent.

Example 19 *Procedure 17 applied to the difference equation*

$$(F_n - 2F_{n+1})u(n+2) + (3F_n + 2F_{n+1})u(n+1) - F_n u(n) = 0$$

with the assumption $\ker \phi = \{0\}$ gives the infinite output

$$\begin{aligned} & F_n^2 \\ & F_n^2(F_n^2 + F_n F_{n+1} - F_{n+1}^2)^2 \\ & F_n^2(F_n^2 + F_n F_{n+1} - F_{n+1}^2)^4 \\ & F_n^2(F_n^2 + F_n F_{n+1} - F_{n+1}^2)^6 \\ & F_n^2(F_n^2 + F_n F_{n+1} - F_{n+1}^2)^8 \\ & \vdots \end{aligned}$$

All these solutions are correct. However, they are not linearly independent as sequences. Indeed, because of the identity $(F_n^2 + F_n F_{n+1} - F_{n+1}^2)^2 = 1$ they all represent the same solution. If we take $\ker \phi = \langle (x_0^2 + x_0 x_1 + x_1^2)^2 - 1 \rangle$, assuming that x_0 and x_1 are the variables encoding F_n and F_{n+1} , respectively, we get the single solution

$$F_n^2$$

as output. There ought to be a second solution to the equation, linearly independent of $u_1(n) = F_n^2$. This second solution is

$$u_2(n) = F_n^2 \sum_{k=1}^{n-1} \frac{1}{F_k^2 F_{k+1}^2} \prod_{i=2}^k \frac{F_{i-1}}{2F_i - F_{i-1}}$$

and cannot be expressed as a rational function in F_n and F_{n+1} . This is the reason why only one solution is output by Procedure 17. (Note, however, that u_2 is an admissible sequence according to Ex. 4.)

6.2 The Inhomogeneous Equation

Extension of Procedure 17 to the inhomogeneous equation (3) is straightforward. If an additional variable z is introduced to represent the inhomogeneous

part $g(n)$, then we have

$$p(f_1(n), \dots, f_m(n)) \text{ solves (3)}$$

$$\iff$$

$$(p(x_{1,0}, \dots, x_{m,0}), \dots, p(x_{1,r}, \dots, x_{m,r}), -1) \in \text{Syz}(y_0, \dots, y_r, z)$$

Modifying lines 3 and 8 of Procedure 17 accordingly, we obtain a method to find the solutions of the inhomogeneous equation.

This leads us to an alternative procedure for indefinite summation of admissible sequences. In order to find a closed form for $\sum_{k=1}^n f(k)$ in terms of some other given admissible sequences $f_1(n), \dots, f_m(n)$, solve the telescoping equation

$$u(n+1) - u(n) = f(n+1)$$

using the inhomogeneous extension of Procedure 17. If $u(n)$ is a solution, then $\sum_{k=1}^n f(k) = u(n) - u(0)$.

Example 20 *For the Legendre polynomials $P_n(x)$ we find the summation identity*

$$\sum_{k=0}^n (2k+1)P_k(x)P_k(y) = \frac{n+1}{x-y}(P_n(y)P_{n+1}(x) - P_n(x)P_{n+1}(y))$$

by solving the telescoping equation

$$u(n+1) - u(n) = (2n+3)P_{n+1}(x)P_{n+1}(y)$$

in terms of $n, P_n(x), P_{n+1}(x), P_n(y), P_{n+1}(y)$.

7 Conclusion

Difference equations of quite complicated form can be solved algorithmically. In this paper, the focus was on quite a large class of univariate sequences that we called admissible. We have given an effective method for enumerating a basis of the ideal of all algebraic dependencies of a set of given admissible sequences, a problem for which a finite algorithm is not likely to be found. Applications related to combinatorial sequences and symbolic summation were indicated.

References

- [1] S.A. Abramov and K.Yu. Kvensenko. Fast algorithms to search for the rational solutions of linear differential equations with polynomial coefficients. In *Proceedings of ISSAC'91*, pages 267–270, 1991.
- [2] Sergei A. Abramov. Rational solutions of linear difference and q -difference equations with polynomial coefficients. In *Proceedings of ISSAC '95*, July 1995.
- [3] Milton Abramowitz and Irene A. Stegun. *Handbook of Mathematical Functions*. Dover Publications, Inc., 9th edition, 1972.
- [4] Andrew Adler. Some recursively unsolvable problems in analysis. *Proceedings of the American Mathematical Society*, 22(2):523–526, 1969.
- [5] Denis Barthou, Paul Feautrier, and Xavier Redon. On the equivalence of two systems of affine recurrence equations. In *Proceedings of EuroPar2002*, pages 1–8, 2002.
- [6] Thomas Becker, Volker Weispfenning, and Heinz Kredel. *Gröbner Bases*. Springer, 1993.
- [7] Alin Bostan, Thomas Cluzeau, and Bruno Salvy. Fast algorithms for polynomial solutions of linear differential equations. In *Proceedings of ISSAC'05*, pages 45–52, 2005.
- [8] F. Boulier, D. Lazard, F. Ollivier, and M. Petitot. Representation for the radical of a finitely generated differential ideal. In *Proceedings of ISSAC'95*, pages 158–166, 1995.
- [9] Manuel Bronstein. On solutions of linear ordinary differential equations in their coefficient field. *Journal of Symbolic Computation*, 13(4):413–439, 1992.
- [10] Bob F. Caviness. On canonical forms and simplification. *Journal of the ACM*, 17(2):385–396, 1970.
- [11] J. Denef and L. Lipshitz. Decision problems for differential equations. *The Journal of Symbolic Logic*, 54(3):941–950, 1989.
- [12] Graham Everest, Alf van der Poorten, Igor Shparlinski, and Thomas Ward. *Recurrence Sequences*, volume 104 of *Mathematical Surveys and Monographs*. American Mathematical Society, 2003.
- [13] David Gale. The strange and surprising saga of the Somos sequences. *Mathematical Intelligencer*, 13(1):40–42, 1991.
- [14] Ronald L. Graham, Donald E. Knuth, and Oren Patashnik. *Concrete Mathematics*. Addison-Wesley, second edition, 1994.
- [15] Vesa Halava, Tero Harju, Mika Hirvensalo, and Juhani Karhumäki. Skolem's problem – on the border between decidability and undecidability, 2005. preprint.

- [16] P.A. Hendriks and M.F. Singer. Solving difference equations in finite terms. *Journal of Symbolic Computation*, 27(3):239–259, 1999.
- [17] Michael Karr. Summation in finite terms. *Journal of the ACM*, 28:305–350, 1981.
- [18] Manuel Kauers. *Algorithms for Nonlinear Higher Order Difference Equations*. PhD thesis, RISC-Linz, Johannes Kepler Universität Linz, 2005.
- [19] Manuel Kauers. SumCracker: A package for manipulating symbolic sums and related objects. *Journal of Symbolic Computation*, 41(9):1039–1057, 2006.
- [20] Manuel Kauers. An algorithm for deciding zero equivalence of nested polynomially recurrent sequences. *Transactions on Algorithms*, 3(2), 2007. Article no. 18.
- [21] Manuel Kauers and Burkhard Zimmermann. Computing the algebraic relations of C-finite sequences and multisequences. *Journal of Symbolic Computation*, 2007. to appear.
- [22] E. R. Kolchin. *Differential Algebra and Algebraic Groups*. Academic Press, 1973.
- [23] Jerald J. Kovacic. An algorithm for solving second order linear homogenous differential equations. *Journal of Symbolic Computation*, 2:3–43, 1986.
- [24] Louis M. Milne-Thomson. *The Calculus of Finite Differences*. Macmillan and Co., ltd., 1933.
- [25] Marko Petkovšek. Hypergeometric solutions of linear recurrences with polynomial coefficients. *Journal of Symbolic Computation*, 14(2–3):243–264, 1992.
- [26] Stanley Rabinowitz. Problem B-951. *The Fibonacci Quarterly*, 40(1):84, 2003.
- [27] Daniel Richardson. Some undecidable problems involving elementary functions of a real variable. *The Journal of Symbolic Logic*, 33(4):514–520, 1968.
- [28] Carsten Schneider. Solving parameterized linear difference equations in terms of indefinite nested sums and products. *Journal of Difference Equations and Applications*, 11(9):799–821, 2005.
- [29] John Shackell. Zero-equivalence in function fields defined by algebraic differential equations. *Transactions of the American Mathematical Society*, 336(1):151–171, 1993.
- [30] Michael F. Singer. Liouvillian solutions of linear differential equations with liouvillian coefficients. *Journal of Symbolic Computation*, 11(3):251–273, 1991.
- [31] Michael Somos. Problem 1470. *Cruz Mathematicorum*, 15:208, 1989.
- [32] Felix Ulmer. Liouvillian solution of third order differential equations. *Journal of Symbolic Computation*, 36:855–889, 2003.

- [33] Felix Ulmer and Michael F. Singer. Liouvillian and algebraic solutions of second and third order linear differential equations. *Journal of Symbolic Computation*, 16(1):37–73, 1993.
- [34] Joris van der Hoeven. A new zero-test for formal power series. In *Proceedings of ISSAC'02*, 2002.
- [35] Alf van der Poorten. Elliptic curves and continued fractions. *Journal of Integer Sequences*, 8(2):1–19, 2005.
- [36] Mark van Hoeij. Rational solutions of linear difference equations. In *Proceedings of ISSAC'98*, pages 120–123, 1998.
- [37] Doron Zeilberger. A holonomic systems approach to special function identities. *Journal of Computational and Applied Mathematics*, 32:321–368, 1990.
- [38] Burkhard Zimmermann. PhD thesis, RISC-Linz, Johannes Kepler Universität Linz, in preparation.