Equivalence Relations and Partitions

Manuel Kauers

TSW-Homework

1 Introduction

Mathematicians are always interested in studying *relations* among certain objects. From an abstract point of view, elements of a set might be related to each other in an arbitrary complicated way, and it is hard to find interesting properties that describe the behaviour of relations in general. On the other hand, if the attention is focussed to a somewhat restricted kind of relations, things might become easier.

In this paper, we will study a special kind of relations, which are at the same time reflexive, transitive, and symmetric. Relations with these properties will be called *equivalence relations.* We will formally introduce the notion of an equivalence relation and study general properties of these relations.

We assume familiarity with the notions of *set,* in particular with the *element of* predicate $\in$, the cartesian product $\times$, the *subset* predicate $\subseteq$ and the operations *union* $\cup$ and *intersection* $\cap$. We will use infix notation for the binary predicates $\in$ and $\subseteq$ as well as for the the binary function symbols $\times$, $\cup$, and $\cap$. By $\varnothing$, we will denote the *empty set.*

By lowercase letters $x, y, \ldots$ we will denote basic elements, capital letters $A, B, \ldots$ will denote sets, and calligraphic letters $\mathcal{A}, \mathcal{B}, \ldots$ will denote families of sets. The elements of a cartesian product $A \times B$ will be denoted by pairs $\langle a, b\rangle$ $(a \in A, b \in B)$.

2 Relations

We start by defining the notion of a relation.

Def. 2.1

1. *Let S be a set and R be a subset of $S \times S$. Then R is called a* relation *on S, formally*

$$\underset{S,R}{\forall}\Big(\text{isRelation}[R,S] \leftrightarrow R \subseteq S \times S\Big).$$

   If R is a relation on S then S is called a base set *of R.*

2. *More general, R is called a* relation, *if R is a relation on some set S, i.e.*

$$\underset{R}{\forall}\Big(\text{isRelation}[R] \leftrightarrow \underset{S}{\exists}\,\text{isRelation}[R,S]\Big).$$

3. *If R is a relation, then the smallest base set of R is called the* domain *of R, denoted by* $\mathrm{dom}[R]$*:*

$$\mathrm{dom}[R] := \{x : \underset{\langle x,y\rangle}{\exists} \langle x,y\rangle \in R \vee \langle x,y\rangle \in R\}.$$

If R is a relation on some set S, we will write aRb as a shortcut of $\langle a,b\rangle \in R$, and we may say "R relates a to b" if aRb is true.

Ex. 2.1 *Let* $S = \{1,2,3,4\}$ *and* $R = \{\langle 1,2\rangle, \langle 2,3\rangle, \langle 3,1\rangle\}$.

- *R is a relation on S.*
- *R is a relation, since R is a relation on S.*
- $\mathrm{dom}[R] = \{1,2,3\}$.
- $\mathrm{dom}[\varnothing] = \varnothing$.

We proceed to consider relations that have special properties.

Def. 2.2

1. *Let S be a set and R a relation on S. R is called* symmetric *on S, if xRy implies yRx for all $x, y \in S$, formally*

$$\underset{\substack{S,R\\ \mathrm{isRelation}[R,S]}}{\forall} \Big(\mathrm{isSymm}[R,S] \leftrightarrow \underset{x,y\in S}{\forall} (xRy \rightarrow yRx) \Big).$$

2. *A relation R is called* symmetric, *if xRy implies yRx for all x, y, formally*

$$\underset{\substack{R\\ \mathrm{isRelation}[R]}}{\forall} \Big(\mathrm{isSymm}[R] \leftrightarrow \underset{x,y}{\forall} (xRy \rightarrow yRx) \Big).$$

In fact, it is easy to prove that these two notions of symmetry are equivalent in the sense that a relation is symmetric if and only if it is symmetric on all of its base sets.

Prop. 2.1

$$\underset{R}{\forall} \Big(\mathrm{isSymm}[R] \leftrightarrow \underset{\substack{S\\ \mathrm{isRelation}[R,S]}}{\forall} \mathrm{isSymm}[R,S] \Big).$$

Proof. Let R be an arbitrary relation. (If R is not a relation, the scope of the first quantifier amounts to false $\leftrightarrow$ false, i.e. it is nothing to prove.)

"$\rightarrow$" Assume that R is symmetric, i.e. assume

$$\underset{x,y}{\forall} (xRy \rightarrow yRx). \tag{A1}$$

by Def. 2.2.2. We have to show that R is symmetric for each base set S, i.e.

$$\underset{\substack{S\\ \mathrm{isRelation}[R,S]}}{\forall} \mathrm{isSymm}[R,S]. \tag{G1}$$

So let S be an arbitrary but fixed base set of R. Then, using Def. 2.2.1, it remains to show

$$\underset{x,y\in S}{\forall}(xRy \to yRx). \tag{G2}$$

For arbitrary $x, y \in S$ this statement is already asserted by A1.

"$\leftarrow$" Assume that R is symmetric for all base sets S, i.e. assume

$$\underset{\substack{S\\ \text{isRelation}[R,S]}}{\forall} \text{isSymm}[R,S]. \tag{A2}$$

We have to show

$$\underset{x,y}{\forall}(xRy \to yRx). \tag{G3}$$

This means, for some arbitrary but fixed x, y with

$$xRy, \tag{A3}$$

it is to show that

$$yRx. \tag{G4}$$

By Def. 2.1.3 and A3, we know that

$$\langle x, y\rangle \in \text{dom}[R]. \tag{A4}$$

Taking dom$[R]$ as S in A2 and using A3 again, we obtain yRx which matches G4.

This completes the proof. ∎

Def. 2.3

1. *Let S be a set and R be a relation on S. R is called* transitive *over S if xRy and yRz implies xRz for all $x, y, z \in S$. Formally,*

$$\underset{\substack{R,S\\ \text{isRelation}[R,S]}}{\forall} \Big(\text{isTrans}[R,S] \leftrightarrow \underset{x,y,z\in S}{\forall}(xRy \wedge yRz \to xRz)\Big)$$

2. *Let R be a relation. R is called* transitive, *if xRy and yRz implies xRz for all x, y, z. Formally,*

$$\underset{\substack{R\\ \text{isRelation}[R]}}{\forall} \Big(\text{isTrans}[R] \leftrightarrow \underset{x,y,z}{\forall}(xRy \wedge yRz \to xRz)\Big)$$

Again, the two notions of transitivity are strongly related.

Prop. 2.2 *(without proof)*

$$\underset{R}{\forall}\Big(\text{isTrans}[R] \leftrightarrow \underset{S}{\forall}\text{isTrans}[R,S]\Big).$$

The proof of Prop. 2.2 is similar to the proof of Prop. 2.1 and left to the reader.

Def. 2.4 *Let S be a set and R a relation on S. R is called* reflexive, *if it relates each element of S to itself, i.e.*

$$\underset{\substack{S,R \\ \text{isRelation}[R,S]}}{\forall} \Big(\text{isRefl}[R,S] \leftrightarrow \underset{x \in S}{\forall} xRx \Big).$$

Note that reflexitivity is only defined with respect to a base set, but not in general. In fact, no general relation R will ever have the property

$$\underset{\substack{S \\ \text{isRelation}[R,S]}}{\forall} \text{isRefl}[R,S], \qquad (*)$$

which would be desired for a general definition in analogy to Prop. 2.1 and Prop. 2.2. The argument is as follows.

Assume that R is a relation with the property $(*)$ and consider an arbitrary base set S of R. Then, $R \subseteq S \times S$ by Def. 2.1.1. By an axiom of set theory, there must be some $x \notin S$. For such an x, consider $S' := S \cup \{x\}$. Then R is also a relation on S' by Def. 2.1.1 and the fact that $A \subseteq B$ implies $A \times A \subseteq B \times B$. But R is not reflexive on S' in contradiction to $(*)$.

This justifies that there is no serious way to define reflexivity without refering to a particular base set. As opposed to transitivity and symmetry, which are properties of the relation itself, reflexivity depends on the relation as well as on its base set.

At the end of this section, we provide the definition of an equivalence relation.

Def. 2.5 *Let $\sim$ be a relation on S. $\sim$ is called an* equivalence relation *on S, if $\sim$ is reflexive, transitive, and symmetric:*

$$\underset{\substack{\sim,S \\ \text{isRelation}[\sim,S]}}{\forall} \Big(\text{isEqRel}[\sim,S] \leftrightarrow (\text{isRefl}[\sim,S] \wedge \text{isSymm}[\sim] \wedge \text{isTrans}[\sim]) \Big)$$

Note that being an equivalence relation is also a property which depends on the base set since reflexivity depends on the base set.

Ex. 2.2 *Let $S = \{1,2,3\}$, and $R = \{\langle 1,1\rangle, \langle 1,2\rangle, \langle 2,1\rangle, \langle 2,2\rangle, \langle 3,3\rangle\}$. Then R is an equivalence relation on S.*

3 Partitions

We denote by $\mathcal{P}[S]$ the *powerset* of a set S. Recall that

$$\underset{S,T}{\forall} \Big(T \in \mathcal{P}[S] \leftrightarrow T \subseteq S \Big)$$

is the defining property of the powerset, i.e. the powerset of S is the set of all subsets of S.

Def. 3.1 *A set $\mathcal{F}$ is called a* family *if $\mathcal{F} \subseteq \mathcal{P}[S]$ for some set S.*

$$\underset{\mathcal{F}}{\forall} \Big(\text{isFamily}[\mathcal{F}] \leftrightarrow \underset{S}{\exists}\, \mathcal{F} \subseteq \mathcal{P}[S] \Big).$$

Hence, a family is a set of sets. We proceed by considering families with special properties.

Def. 3.2 *Let $\mathcal{F}$ be a family.*

1. *$\mathcal{F}$ is called a* cover *of a set S if every element $x \in S$ is contained in at least one of the elements of $\mathcal{F}$:*

$$\underset{\substack{\mathcal{F},S \\ \text{isFamily}[\mathcal{F}]}}{\forall} \Big(\text{isCover}[\mathcal{F}, S] \leftrightarrow \underset{x \in S}{\forall} \underset{T \in \mathcal{F}}{\exists} x \in T \Big)$$

2. *$\mathcal{F}$ is called* disjoint *if two distinct elements of $\mathcal{F}$ don't have any element in common:*

$$\underset{\substack{\mathcal{F} \\ \text{isFamily}[\mathcal{F}]}}{\forall} \Big(\text{isDisj}[\mathcal{F}] \leftrightarrow \underset{P,Q \in \mathcal{F}}{\forall} (P \neq Q \rightarrow P \cap Q = \varnothing) \Big)$$

3. *$\mathcal{F}$ is called a* partition *of a set S if $\mathcal{F}$ is a disjoint cover of S:*

$$\underset{\substack{\mathcal{F},S \\ \text{isFamily}[\mathcal{F}]}}{\forall} \Big(\text{isPartition}[\mathcal{F}, S] \leftrightarrow \text{isCover}[\mathcal{F}, S] \wedge \text{isDisjoint}[\mathcal{F}] \Big)$$

Ex. 3.1 *Let $\mathcal{F}_1 = \{\{1,2\},\{2,3\},\{3,4\}\}$, $\mathcal{F}_2 = \{\{1,2\},\{3,4\}\}$.*

- *Both $\mathcal{F}_1$ and $\mathcal{F}_2$ are covers of $\{1,2,3,4\}$ but not of $\{1,2,3,4,5\}$.*
- *Each family covers the empty set.*
- *$\mathcal{F}_2$ is disjoint, $\mathcal{F}_1$ is not.*
- *$\mathcal{F}_2$ is a partition of $\{1,2,3,4\}$*
- *$\{S\}$ is always a partition of S.*

4 Equivalence Relations and Partitions

This section links the notions of equivalence relation (Def. 2.5) and of partition (Def. 3.2.3) to each other. In fact, we will see that each equivalence relation on a set S defines a partition of S, and vice versa.

Def. 4.1 *Let S be a set, $\sim$ an equivalence relation on S, and $x \in S$. Then, the set*

$$[x]_\sim := \{y : y \in S \wedge y \sim x\}$$

is called the equivalence class *of x.*

Prop. 4.1 *Let S be a set, $\sim$ an equivalence relation on S, and $x, y \in S$ with $x \sim y$. Then $[x]_\sim = [y]_\sim$.*

$$\underset{\substack{S,\sim \\ \text{isEqRel}[\sim,S]}}{\forall} \underset{x,y \in S}{\forall} (x \sim y \to [x]_\sim = [y]_\sim)$$

Proof. Let S be a set and $\sim$ be an equivalence relation on S, i.e. assume

$$\text{isEqRel}[\sim, S] \tag{A1}$$

which by Def. 2.5 is equivalent to

$$\text{isRefl}[\sim, S] \wedge \text{isSymm}[\sim] \wedge \text{isTrans}[\sim] \tag{A2}$$

which in turn yields

$$\underset{x \in S}{\forall} x \sim x, \tag{A3}$$

$$\underset{x,y}{\forall} (x \sim y \to y \sim x), \tag{A4}$$

$$\underset{x,y,z}{\forall} (x \sim y \wedge y \sim z \to x \sim z) \tag{A5}$$

by definitions 2.4, 2.2.2, and 2.3.2. We have to show

$$\underset{x,y \in S}{\forall} (x \sim y \to [x]_\sim = [y]_\sim) \tag{G1}$$

so consider some arbitrary but fixed $x, y \in S$ with the property

$$x \sim y \tag{A6}$$

and show

$$[x]_\sim = [y]_\sim. \tag{G2}$$

We proceed to show that the left hand side is contained in the right hand side and vice versa.

- We show

  $$[x]_\sim \subseteq [y]_\sim \tag{G3}$$

  which by the definition of $\subseteq$ means that

  $$\underset{z}{\forall} (z \in [x]_\sim \to z \in [y]_\sim) \tag{G4}$$

  Consider an arbitrary but fixed

  $$z \in [x]_\sim \tag{A7}$$

  and show

  $$z \in [y]_\sim. \tag{G5}$$

  By A7 and Def. 4.1, we know that

  $$z \sim x \tag{A8}$$

  and using A8, A6 and A5 we obtain

  $$z \sim y. \tag{A9}$$

  Now G5 follows from A9 and the definition of $[y]_\sim$.

- As for the other direction, we have to show

$$[y]_\sim \subseteq [x]_\sim \tag{G6}$$

which means

$$\underset{z}{\forall}(z \in [y]_\sim \to z \in [x]_\sim) \tag{G7}$$

Take an arbitrary but fixed

$$z \in [y]_\sim \tag{A10}$$

and show

$$z \in [x]_\sim. \tag{G8}$$

By A10 and Def. 4.1, we know that

$$z \sim y \tag{A11}$$

and using A4 and A6 it follows that

$$y \sim x. \tag{A12}$$

By A11, A12, and A5 we get

$$z \sim x. \tag{A13}$$

Now G8 follows from A13 and the definition of $[x]_\sim$.

This completes the proof. ∎

Theorem 4.1 *Let S be a set, $\sim$ an equivalence relation on S, and let*

$$\mathcal{F} = \{[x]_\sim : x \in S\}$$

Then $\mathcal{F}$ is a partition:

$$\underset{\substack{S,\sim \\ \text{isEqRel}[\sim,S]}}{\forall} \text{isPartition}[\{[x]_\sim : x \in S\}, S].$$

Proof. Let S be a set and $\sim$ be an equivalence relation on S, i.e. assume

$$\text{isEqRel}[\sim, S] \tag{A1}$$

and show

$$\text{isPartition}[\{[x]_\sim : x \in S\}, S]. \tag{G1}$$

By Def. 2.5, A1 leads to

$$\text{isRefl}[\sim, S] \wedge \text{isSymm}[\sim] \wedge \text{isTrans}[\sim] \tag{A2}$$

which might be split and expanded into

$$\underset{x \in S}{\forall} x \sim x, \tag{A3}$$

$$\underset{x,y}{\forall}(x \sim y \to y \sim x), \tag{A4}$$

$$\underset{x,y,z}{\forall}(x \sim y \wedge y \sim z \to x \sim z) \tag{A5}$$

using the definitions 2.4, 2.2.2, and 2.3.2. Writing $\mathcal{F} = \{[x]_\sim : x \in S\}$ and using Def. 3.2.3 we can rewrite G1 as

$$\text{isCover}[\mathcal{F}, S] \wedge \text{isDisjoint}[\mathcal{F}] \tag{G2}$$

i.e. we have to show that $\mathcal{F}$ is a disjoint cover of S.

- We first show

$$\text{isCover}[\mathcal{F}, S] \tag{G3}$$

which by Def. 3.2.1 means

$$\underset{x\in S}{\forall}\ \underset{T\in\mathcal{F}}{\exists}\ x \in T. \tag{G4}$$

Take an arbitrary but fixed $x \in S$ and show

$$\underset{T\in\mathcal{F}}{\exists}\ x \in T. \tag{G5}$$

We claim that $T = [x]_\sim$ does the job, i.e. we claim that

$$x \in [x]_\sim. \tag{G6}$$

By Def. 4.1 we know that

$$\underset{y\in S}{\forall}\ y \in [x]_\sim \leftrightarrow y \sim x. \tag{A6}$$

Taking $y = x$ we can conlcude

$$x \in [x]_\sim \leftrightarrow x \sim x. \tag{A7}$$

With the aid of A7, G6 might be rewritten as

$$x \sim x. \tag{G7}$$

Since $x \in S$, G7 is asserted by A3.

- Now we prove

$$\text{isDisjoint}[\mathcal{F}], \tag{G8}$$

which by Def. 3.2.2 means

$$\underset{P,Q\in\mathcal{F}}{\forall} P \neq Q \rightarrow P \cap Q = \varnothing. \tag{G9}$$

For arbitrary $[x]_\sim, [y]_\sim \in F$ with

$$[x]_\sim \neq [y]_\sim \tag{A8}$$

we will prove

$$[x]_\sim \cap [y]_\sim = \varnothing. \tag{G10}$$

Assume the contrary of G10, i.e. the existence of an element z with

$$z \in [x]_\sim \cap [y]_\sim \tag{A9}$$

which means

$$z \in [x]_\sim \tag{A10}$$
$$z \in [y]_\sim. \tag{A11}$$

By Def. 4.1, this can be written as

$$z \sim x \tag{A12}$$
$$z \sim y. \tag{A13}$$

By A4 we can transform A13 into

$$y \sim z. \tag{A14}$$

and A5 allows to conclude

$$x \sim y \tag{A15}$$

from A12 and A15. Using Prop. 4.1, we arrive at

$$[x]_\sim = [y]_\sim \tag{A16}$$

which contradicts A8. Hence assumption A9 was wrong and therefore G10 must be true.

This completes the proof. ∎

The last theorem showed that an equivalence relation on a set S defines a partition on S. We will now turn to the "reverse" statement, namely that every partition of S also defines an equivalence relation on S. This requires a little more preparation.

Prop. 4.2 *Let S be a set and $\mathcal{F}$ be a partition of S. Then, for each $x \in S$ there exists exactly one $P \in \mathcal{F}$ such that $x \in P$, i.e.*

$$\underset{\substack{S,\mathcal{F} \\ \text{isPartition}[\mathcal{F},S]}}{\forall} \Big(\underset{x \in S}{\forall} \big(\underset{P \in \mathcal{F}}{\exists} (x \in P) \wedge \underset{P,Q \in \mathcal{F}}{\forall} (x \in P \wedge x \in Q \to P = Q) \big) \Big).$$

Proof. Let S be an arbitrary set and $\mathcal{F}$ be an arbitrary partition of S, i.e. assume

$$\text{isPartition}[\mathcal{F}, S]. \tag{A1}$$

We have to show

$$\underset{x \in S}{\forall} \big(\underset{P \in \mathcal{F}}{\exists} (x \in P) \wedge \underset{P,Q \in \mathcal{F}}{\forall} (x \in P \wedge x \in Q \to P = Q) \big). \tag{G1}$$

First we expand the definition of partition in A1. By Def. 3.2.3 this leads to

$$\text{isDisjoint}[\mathcal{F}], \tag{A2}$$
$$\text{isCover}[\mathcal{F}, S], \tag{A3}$$

which in turn by Def. 3.2.1 and Def. 3.2.2 lead to

$$\underset{x \in S}{\forall} \underset{T \in F}{\exists} x \in T \tag{A4}$$
$$\underset{P,Q \in \mathcal{F}}{\forall} (P \neq Q \to P \cap Q = \varnothing). \tag{A5}$$

Now let $x \in S$ be arbitrary but fixed. We have to show

$$\underset{P \in \mathcal{F}}{\exists} x \in P \tag{G2}$$
$$\underset{P,Q \in \mathcal{F}}{\forall} (x \in P \wedge x \in Q \to P = Q). \tag{G3}$$

G2 is immediate by assumption A4. As for G3, consider arbitrary but fixed $P, Q \in F$ and assume

$$x \in P \wedge x \in Q, \tag{A6}$$

which is equivalent to $x \in P \cap Q$. It remains to show that

$$P = Q. \tag{G4}$$

By A5, we know

$$P \neq Q \to P \cap Q = \varnothing \tag{A7}$$

Assuming that G4 does not hold, A7 leads to the conclusion $P \cap Q = \varnothing$ in contradiction to A6. Thus, G4 holds which completes the proof. ∎

The fact that every element $x \in S$ is contained in exactly one element of a partition $\mathcal{F}$ of S shows that the following notion is well defined.

Def. 4.2 *Let S be a set, $\mathcal{F}$ a partition of S, and $x \in S$. The* home *of x with respect to $\mathcal{F}$, denoted by* $\mathrm{home}[x, \mathcal{F}]$*, is defined as the uniquely determined element $P \in \mathcal{F}$ with $x \in P$:*

$$\underset{\substack{S,\mathcal{F} \\ \mathrm{isPartition}[\mathcal{F},S]}}{\forall} \Big(\underset{x \in S}{\forall} \underset{P \in \mathcal{F}}{\forall} P = \mathrm{home}[x, \mathcal{F}] \leftrightarrow x \in P \Big).$$

Theorem 4.2 *Let S be a set and $\mathcal{F}$ be a partition of S. Furthermore, let*

$$\sim \; := \{\langle x, y \rangle : x, y \in S \wedge \mathrm{home}[x, \mathcal{F}] = \mathrm{home}[y, \mathcal{F}]\}.$$

Then $\sim$ is an equivalence relation.

$$\underset{\substack{S,\mathcal{F} \\ \mathrm{isPartition}[\mathcal{F},S]}}{\forall} \mathrm{isEqRel}[\{\langle x, y \rangle : x, y \in S \wedge \mathrm{home}[x, \mathcal{F}] = \mathrm{home}[y, \mathcal{F}]\}, S].$$

Proof. Let S be a set and $\mathcal{F}$ be a partition of S, i.e. assume

$$\mathrm{isPartition}[\mathcal{F}, S] \tag{A1}$$

which by Def. 3.2.3 means

$$\mathrm{isDisj}[\mathcal{F}] \tag{A2}$$

$$\mathrm{isCover}[\mathcal{F}, S] \tag{A3}$$

which by Def. 3.2.1 and Def. 3.2.2 means

$$\underset{P,Q \in \mathcal{F}}{\forall} P \neq Q \to P \cap Q = \varnothing \tag{A4}$$

$$\underset{x \in S}{\forall} \underset{P \in \mathcal{F}}{\exists} x \in P. \tag{A5}$$

We have to prove

$$\mathrm{isEqRel}[\{\langle x, y \rangle : x, y \in S \wedge \mathrm{home}[x, \mathcal{F}] = \mathrm{home}[y, \mathcal{F}]\}, S]. \tag{G1}$$

Using the abbreviation

$$R := \{\langle x, y \rangle : x, y \in S \wedge \mathrm{home}[x, \mathcal{F}] = \mathrm{home}[y, \mathcal{F}]\} \tag{A6}$$

and Def. 2.5, G1 is equivalent to

$$\mathrm{isRefl}[R, S] \wedge \mathrm{isSymm}[R] \wedge \mathrm{isTrans}[R]. \tag{G2}$$

We prove the three properties separately.

- Reflexivity. We prove

$$\text{isRefl}[R, S] \tag{G3}$$

which means

$$\underset{x \in S}{\forall} xRx \tag{G4}$$

by Def. 2.4. Let $x \in S$ be arbitrary but fixed and prove

$$xRx, \tag{G5}$$

which by A6 is equivalent to

$$\text{home}[x, \mathcal{F}] = \text{home}[x, \mathcal{F}] \tag{G6}$$

which is obviously the case.

- Symmetry. We prove

$$\text{isSymm}[R] \tag{G7}$$

which means

$$\underset{x,y}{\forall} xRy \to yRx. \tag{G8}$$

Now take x, y arbitrary, assume

$$xRy \tag{A7}$$

and show

$$yRx. \tag{G9}$$

By the definition of R (A6), A7 is equivalent to

$$\text{home}[x, \mathcal{F}] = \text{home}[y, \mathcal{F}] \tag{A8}$$

and G9 is equivalent to

$$\text{home}[y, \mathcal{F}] = \text{home}[x, \mathcal{F}]. \tag{G10}$$

G10 is obtained by A8 using the symmetry of "=."

- Transitivity. We prove

$$\text{isTrans}[R] \tag{G11}$$

which means

$$\underset{x,y,z}{\forall} xRy \wedge yRz \to xRz. \tag{G12}$$

Now take x, y, z arbitrary, assume

$$xRy, \tag{A9}$$
$$yRz, \tag{A10}$$

and prove

$$xRz. \tag{G13}$$

By the definition of R (A6), A9 and A10 yield the equivalent statements

$$\text{home}[x, \mathcal{F}] = \text{home}[y, \mathcal{F}], \tag{A11}$$
$$\text{home}[y, \mathcal{F}] = \text{home}[z, \mathcal{F}], \tag{A12}$$

while

$$\text{home}[x, \mathcal{F}] = \text{home}[z, \mathcal{F}] \tag{G14}$$

remains to be proven. A11 and A12 together with the transitivity of "=" give G14.

Finally, we have shown that R is reflexive, symmetric and transitive, thus R is an equivalence relation. ∎

5 Conclusion

We wasted 12 pages of paper and several hours of work to prove the trivial fact that equivalence relations correspond to partitions, which is usually left as an exercise to first year students. Compared to an informal proof of a few lines as might be given by the first year student, we gained no additional insight besights the fact that it "really" works as we expected. The material provided is not worth the paper on which it is written.