

An Explicit Expression of the Lüroth Invariant

Romain Basson
IRMAR
Université de Rennes 1,
Campus de Beaulieu
35042 Rennes
France
romain.basson@univ-
rennes1.fr

Christophe Ritzenthaler
Institut de Mathématiques de
Luminy, UMR 6206 du CNRS,
Luminy, Case 907
13288 Marseille
France
ritzenth@iml.univ-mrs.fr

Reynald Lercier
DGA & IRMAR
Université de Rennes 1,
Campus de Beaulieu
35042 Rennes
France
reynald.lercier@m4x.org

Jeroen Sijsling
Mathematics Institute,
Zeeman Building, University of
Warwick
Coventry CV4 7AL
United Kingdom
sijsling@gmail.com

ABSTRACT

In this short note, we give an algorithm that returns an explicit expression of the Lüroth invariant in terms of the Dixmier-Ohno invariants of plane quartic curves. We also obtain an explicit factorized expression on the locus of Ciani quartics in terms of the coefficients. After this calculation, we extend our methods to answer two open theoretical questions concerning the sub-locus of singular Lüroth quartics.

Categories and Subject Descriptors

J.2 [Computer applications]: Mathematics and statistics

Keywords

Invariant ; ternary quartic ; genus 3 ; Ciani quartic ; algorithm ; Dixmier-Ohno invariants.

1. LÜROTH QUARTICS

This note considers *Lüroth quartics*, which are plane quartics containing the ten vertices of a non-degenerate pentilateral. To make these notions precise, we give the following definitions. Let V be a three-dimensional vector space over the complex field \mathbb{C} .

DEFINITION 1. *A complete pentilateral in the projective plane $\mathbb{P}V$ is a curve $C \subset \mathbb{P}V$ consisting of the union of five lines ℓ_1, \dots, ℓ_5 that are three by three linearly independent (which is to say that the pairwise intersections of the lines ℓ_i yield exactly 10 distinct points).*

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.

ISSAC'13, June 26–29, 2013, Boston, Massachusetts, USA.
Copyright 2013 ACM 978-1-4503-2059-7/13/06 ...\$15.00.

The vertices of a complete pentilateral C are the double points of C , that is, the 10 points $\bigcup_{i \neq j} ((\ell_i = 0) \cap (\ell_j = 0))$.

DEFINITION 2. *Let $Q \subset \mathbb{P}V$ be a non-singular quartic. Then Q is called a non-singular Lüroth quartic if it contains the vertices of a complete pentilateral in $\mathbb{P}V$.*

The set of plane quartics in $\mathbb{P}V$ can be identified with the projective space $\mathbb{P}\text{Sym}^4(V^*)$ over the fourth symmetric power of the dual vector space V^* of V . This inherits an action of the group $\text{GL}(V)$ and its derived subgroup $\text{SL}(V)$, which act canonically on V . Choosing a basis, as we will do during our calculations, identifies V with \mathbb{C}^3 and the set of quartics $\mathbb{P}\text{Sym}^4(V^*)$ with the projectivization of the vector space on the homogeneous degree 4 monomials in the canonical basis x, y, z of the dual space $(\mathbb{C}^3)^*$. We will in turn identify this projective space with \mathbb{P}^{14} by choosing some ordering of these 15 monomials. In this way, \mathbb{P}^{14} inherits an action of the groups $\text{GL}_3(\mathbb{C})$ and $\text{SL}_3(\mathbb{C})$.

The classical study of Lüroth quartics culminated in 1919 with the work of Morley [9]. This showed that the Zariski closure of the locus of non-singular Lüroth quartics in the projective space $\mathbb{P}\text{Sym}^4(V^*)$ is an irreducible hypersurface described as the vanishing locus of a single homogeneous polynomial function L on the projective space of quartics $\mathbb{P}\text{Sym}^4(V^*)$, well-defined up to scalars. We shall call this L the *Lüroth invariant*. Morley showed that L is of degree 54.

DEFINITION 3. *Let $Q \subset \mathbb{P}V$ be a quartic. Then Q is called a Lüroth quartic if $L(Q) = 0$.*

In recent years, after the seminal work of [1], several authors have revived this subject in [2, 11, 12, 13] (see also [14] on the undulation invariant).

However, an explicit expression of L was still missing. In the following section we explain how to compute such an expression. Our main new technique lies in an effective use of [10], an unfortunately unpublished article in which Ohno gives a complete set of generators for the invariants of ternary quartics under the action of $\text{SL}_3(\mathbb{C})$, completing the set of primary invariants found in [3]. These invariants were

also used in [5], and new effective methods to verify their correctness can be found in [4]. Our calculations use the implementation of these invariants in `Magma`, which is due to Kohel.

2. THE ALGORITHM

The key point is the following observation:

PROPOSITION 1. *The homogeneous polynomial function L on $\mathbb{P}\mathrm{Sym}^4(V^*)$ is $\mathrm{GL}(V)$ -invariant up to scalars. In particular, L is $\mathrm{SL}(V)$ -invariant.*

PROOF. Since we are working over an algebraically closed field, this is obvious from the obvious fact that any $\mathrm{GL}(V)$ -transform of a Lüroth quartic is again a Lüroth quartic. \square

Let

$$R = S[\mathbb{P}\mathrm{Sym}^4(V^*)]^{\mathrm{SL}(V)}$$

be the ring of $\mathrm{SL}(V)$ -invariant homogeneous polynomial functions on $\mathbb{P}\mathrm{Sym}^4(V^*)$, which coincides with those functions that are $\mathrm{GL}(V)$ -invariant up to scalars. The structure of the ring R is known. Indeed, let $I = (I_3, I_6, I_9, I_{12}, I_{15}, I_{18}, I_{27})$ be the primary invariants for ternary quartics under the action of $\mathrm{SL}_3(\mathbb{C})$ found by Dixmier in [3], and let $J = (J_9, J_{12}, J_{15}, J_{18}, J_{21}, J_{27})$ be the secondary invariants found by Ohno [10]. In both cases, the index specifies the degree of the invariant as a homogeneous function. Then we have:

THEOREM 1 (DIXMIER-OHNO). *We have $R = \mathbb{C}[I, J]$.*

We now choose a basis for V and a corresponding coordinate system of $\mathbb{P}\mathrm{Sym}^4(V^*) \cong \mathbb{P}^{14}$ as the projective space over the degree 4 monomials in x, y, z . The function L then becomes a homogeneous expression in the coefficients of these monomials. It is unlikely that this function can be written down in any reasonable way (see also the final remark in Section 5). However, by Proposition 1 and Theorem 1, we can express L as a polynomial in the invariant functions in I and J . This expression is not unique, since as we shall see, there are relations between these invariant monomials in degree 54.

To obtain an expression for L , we apply the method of evaluation-interpolation. This is based on the following observation:

PROPOSITION 2. *Let $S = \mathbb{C}[x_3, \dots, x_{27}, y_9, \dots, y_{21}, y'_{21}]$ be a graded polynomial algebra in 13 variables, weighted by indices, and consider the surjection q given by*

$$\begin{aligned} q : S &\longrightarrow R \\ x_k &\longmapsto I_k \\ y_\ell &\longmapsto J_\ell. \end{aligned}$$

Let $R_{54} \subset R$ be the set of homogeneous functions of degree 54, and define $S_{54} \subset S$ analogously. Let K be the kernel of the map $S_{54} \rightarrow R_{54}$. Then $\dim(K) = 215$.

Let X be a finite set of Lüroth quartics. Consider the linear map $q' : S_{54} \rightarrow \mathbb{C}^X$ given by evaluating at the polynomials in X . Let K' be the kernel of q' , and suppose that $\dim(K') = 216$. Let \tilde{L} be an element of $K' \setminus K$. Then the image $L = q(\tilde{L})$ equals the Lüroth invariant.

PROOF. One calculates that the $\dim(S_{54}) = 1380$. Since calculating the Hilbert polynomial of R as in [15, p.1045]

yields $\dim(S_{54}) = 1165$, we indeed find that $\dim(K) = 215$. The rest is straightforward considering the uniqueness of L up to scalars. \square

The details of the calculation are therefore as follows.

1. Construct the 1380 monomials

$$\mathcal{I} = \{I_3^{18}, I_3^{16} I_6, I_3^{15} I_9, I_3^{15} J_9, \dots, J_{18}^3, I_{27}^2\}$$

of degree 54 that generate the \mathbb{C} -vector space of invariants of degree 54.

2. Generate a sufficiently large finite set \mathcal{Q} of cardinality q of random plane quartics with rational coefficients.
3. Generate a sufficiently large finite set \mathcal{L} of cardinality l of random Lüroth quartics of the form

$$\begin{aligned} &\ell_1 \ell_2 \ell_3 \ell_4 + c_1 \cdot \ell_2 \ell_3 \ell_4 \ell_5 + c_2 \cdot \ell_1 \ell_3 \ell_4 \ell_5 \\ &+ c_3 \cdot \ell_1 \ell_2 \ell_4 \ell_5 + c_4 \cdot \ell_1 \ell_2 \ell_3 \ell_5 \end{aligned}$$

where $\ell_1 = x, \ell_2 = y, \ell_3 = z, \ell_4 = x + y + z, \ell_5$ is a line with random rational coefficients and c_i are rational coefficients.

4. Compute the matrix $M_1 = (I(q))_{I \in \mathcal{I}, q \in \mathcal{Q}}$, evaluating the monomials in \mathcal{I} at the quartics in \mathcal{Q} .
5. Compute the matrix $M_2 = (I(q))_{I \in \mathcal{I}, q \in \mathcal{L}}$, evaluating the monomials in \mathcal{I} at the Lüroth quartics in \mathcal{L} .
6. Compute the 215-dimensional kernel N_1 of M_1 . This gives a basis of the homogeneous relations of degree 54 that are satisfied by the invariants of all ternary quartics.
7. Compute the 216-dimensional kernel N_2 of M_2 . This gives a basis of the homogeneous relations of degree 54 that are satisfied by all Lüroth quartics.
8. A non-zero element in the complement of N_2 in N_1 is an expression for L in terms of the Dixmier-Ohno invariants.

All these computations were done with `Magma` software. Over finite fields \mathbb{F}_p with prime cardinality $p = 2017, 10007, 100003$ or even 1000003, computations can be done in less than a minute. However, getting the result over the rationals is more challenging. The main concern is to deal with matrices M_1 and M_2 whose coefficients are as small as possible. So, at Step (2) of the algorithm, we generate plane quartics with random integer coefficients only equal to $-1, 0, 1$. Similarly, we restrict Step (3) to Lüroth forms defined by integer coefficients c_i bounded in absolute value by 4.

We can estimate the size of the computations involved in this run of the algorithm by using the Hadamard bounds for our matrices M_1 and M_2 ; the quartics under consideration yield bounds slightly smaller than 2^{200000} for M_1 and 2^{350000} for M_2 . As a sanity check before running the code over the rationals, we verify that this subset of quartics yields a valid result modulo small primes.

Most of the time is spent at Step (6) and Step (7) of the algorithm, precisely 5 and 9 hours on our laptop (based on a INTEL CORE I7 M620 2.67GHZ processor).

A program to get the result is available on the web page of the authors¹. It uses the implementation of the Dixmier-Ohno invariants in Magma by Kohel². The 1.4Mb result is also available online³. It is given by 1164 monomials with rational coefficients, the largest of which is a quotient of a 680-digit integer by a coprime 671-digit integer. Modulo 1000003, the expression starts as

$$\begin{aligned}
& I_3^{18} + 469313I_3^2I_6^8 + 710780I_6^9 + 969230I_3^3I_6^6I_9 \\
& + 374233I_3I_6^7I_9 + 276144I_3^2I_6^5I_9^2 \\
& + 602674I_6^6I_9^2 + 527614I_3^3I_6^3I_9^3 \\
& + 538637I_3I_6^4I_9^3 + 392526I_3^4I_6I_9^4 \\
& + 645841I_3^2I_6^4I_9^4 + 914224I_3^3I_6^4I_9^4 + 207808I_3^3I_9^5 \\
& + 31577I_3I_6I_9^5 + 635768I_9^6 + 668878I_3^{15}J_9 \\
& + 507293I_3^3I_6^6J_9 + 318476I_3I_6^7J_9 \\
& + 59775I_3^2I_6^5I_9J_9 + 581086I_6^6I_9J_9 \\
& + 830307I_3^3I_6^3I_9^2J_9 + 804817I_3I_6^4I_9^2J_9 \\
& + 6418I_3^3I_9^3J_9 + 578316I_3^4I_6I_9^3J_9 \\
& + 741618I_3^2I_6^3I_9^3J_9 + 452974I_6^3I_9^3J_9 \\
& + 36214I_3^3I_9^4J_9 + 522408I_3I_6I_9^4J_9 \\
& + 253043I_9^5J_9 + 469299I_3^2I_6^5J_9^2 + \dots
\end{aligned}$$

3. CIANI QUARTICS

We call a *Ciani quartic* a plane quartic of the form

$$ax^4 + bx^2y^2 + cx^2z^2 + dy^4 + ey^2z^2 + fz^4.$$

A generic Ciani quartic has automorphism group isomorphic with $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, and conversely every quartic with this property is \mathbb{C} -isomorphic to a Ciani quartic. The dimension of the substratum of Ciani quartics in the full dimension 6 moduli space of plane quartics equals 3.

In [7, Sec.5], using different techniques, Hauenstein and Sottile obtained the factorization on Ciani quartics of the Lüroth invariant as

$$G^4 H^2 J$$

with $G, H, J \in \mathbb{C}[a, b, c, d, e, f]$ homogeneous of respective degree 6, 9 and 12. Using our expression, it is easy to confirm their decomposition. We give a slightly different version of the result, which is due to the fact the coefficients b, c, e are replaced by $2b, 2c, 2e$ in [7, Sec.5]:

$$G = a \cdot d \cdot f \cdot (adf - (1/4)ae^2 - (1/4)b^2f - (1/4)bce - (1/4)c^2d),$$

$$\begin{aligned}
H = & (adf - (1/4)ae^2 - (1/4)b^2f + (1/4)bce + (3/4)c^2d) \\
& \cdot (adf - (1/4)ae^2 + (3/4)b^2f + (1/4)bce - (1/4)c^2d) \\
& \cdot (adf + (3/4)ae^2 - (1/4)b^2f + (1/4)bce - (1/4)c^2d),
\end{aligned}$$

¹<http://iml.univ-mrs.fr/~ritzenth/programme/luroth/luroth.m>

²<http://echidna.maths.usyd.edu.au/kohel/alg/index.html>

³<http://iml.univ-mrs.fr/~ritzenth/programme/luroth/LurothInvF.m>

$$\begin{aligned}
J = & a^4d^4f^4 - (1/49)a^4d^3e^2f^3 + (51/19208)a^4d^2e^4f^2 \\
& - (1/38416)a^4de^6f + (1/614656)a^4e^8 - (1/49)a^3b^2d^3f^4 \\
& - (205/9604)a^3b^2d^2e^2f^3 - (3/38416)a^3b^2de^4f^2 \\
& + (1/153664)a^3b^2e^6f + (15/343)a^3bcd^3ef^3 \\
& + (29/9604)a^3bcd^2e^3f^2 - (5/38416)a^3bcde^5f \\
& - (1/153664)a^3bce^7 - (1/49)a^3c^2d^4f^3 \\
& - (205/9604)a^3c^2d^3e^2f^2 - (3/38416)a^3c^2d^2e^4f \\
& + (1/153664)a^3c^2de^6 + (51/19208)a^2b^4d^2f^4 \\
& - (3/38416)a^2b^4de^2f^3 + (3/307328)a^2b^4e^4f^2 \\
& + (29/9604)a^2b^3cd^2ef^3 - (5/19208)a^2b^3cde^3f^2 \\
& - (3/153664)a^2b^3ce^5f - (205/9604)a^2b^2c^2d^3f^3 \\
& + (2/2401)a^2b^2c^2d^2e^2f^2 + (55/153664)a^2b^2c^2de^4f \\
& + (3/307328)a^2b^2c^2e^6 + (29/9604)a^2bc^3d^3ef^2 \\
& - (5/19208)a^2bc^3d^2e^3f - (3/153664)a^2bc^3de^5 \\
& + (51/19208)a^2c^4d^4f^2 - (3/38416)a^2c^4d^3e^2f \\
& + (3/307328)a^2c^4d^2e^4 - (1/38416)ab^6df^4 \\
& + (1/153664)ab^6e^2f^3 - (5/38416)ab^5cdef^3 \\
& - (3/153664)ab^5ce^3f^2 - (3/38416)ab^4c^2d^2f^3 \\
& + (55/153664)ab^4c^2de^2f^2 + (3/153664)ab^4c^2e^4f \\
& - (5/19208)ab^3c^3d^2ef^2 - (17/76832)ab^3c^3de^3f \\
& - (1/153664)ab^3c^3e^5 - (3/38416)ab^2c^4d^3f^2 \\
& + (55/153664)ab^2c^4d^2e^2f + (3/153664)ab^2c^4de^4 \\
& - (5/38416)abc^5d^3ef - (3/153664)abc^5d^2e^3 \\
& - (1/38416)ac^6d^4f + (1/153664)ac^6d^3e^2 + (1/614656)b^8f^4 \\
& - (1/153664)b^7cef^3 + (1/153664)b^6c^2df^3 \\
& + (3/307328)b^6c^2e^2f^2 - (3/153664)b^5c^3def^2 \\
& - (1/153664)b^5c^3e^3f + (3/307328)b^4c^4d^2f^2 \\
& + (3/153664)b^4c^4de^2f + (1/614656)b^4c^4e^4 \\
& - (3/153664)b^3c^5d^2ef - (1/153664)b^3c^5de^3 \\
& + (1/153664)b^2c^6d^3f + (3/307328)b^2c^6d^2e^2 \\
& - (1/153664)bc^7d^3e + (1/614656)c^8d^4.
\end{aligned}$$

The product G^4H^2J has 1695 monomials. Note that the total amount of weighted monomials in a, b, c, d, e and f in a generic degree 54 invariant is 3439.

4. SINGULAR LÜROTH QUARTICS

Let $\mathcal{L} \subset \mathbb{P}\text{Sym}^4(V^*)$ be the locus of Lüroth quartics, and let $\mathcal{D} \subset \mathbb{P}\text{Sym}^4(V^*)$ be the discriminantal hypersurface defined by the equation $I_{27} = 0$. We will now obtain new results on the geometry of the locus $\mathcal{L} \cap \mathcal{D}$ of *singular Lüroth quartics*. Work by Le Potier and Tikhomirov [8] shows that

$$\mathcal{L} \cap \mathcal{D} = \mathcal{L}_1 \cup \mathcal{L}_2,$$

where \mathcal{L}_1 and \mathcal{L}_2 are irreducible subschemes of $\mathbb{P}\text{Sym}^4(V^*)$ of codimension 2 whose respective degrees as subschemes of \mathcal{D} equal 24 and 30 respectively. Moreover, while \mathcal{L}_1 is reduced, the reduced subscheme $(\mathcal{L}_2)_{\text{red}}$ of \mathcal{L}_2 is of degree 15.

In [12], Ottaviani and Sernesi showed that no new degree 15 invariant vanishes on $(\mathcal{L}_2)_{\text{red}}$, which implies that this scheme is not a principal hypersurface in \mathcal{L} . We will prove a stronger result, namely that none of $\mathcal{L}_1, \mathcal{L}_2, (\mathcal{L}_2)_{\text{red}}$

is a complete intersection. We apply the same methods as in Section 2. The main problem now is to generate quartics in \mathcal{L}_1 and \mathcal{L}_2 .

For \mathcal{L}_2 , we can proceed by using Remark 3.3 in [12]: we now choose the lines in Step (3) above such that three of them have a common point of intersection.

For \mathcal{L}_1 , we have to perform the constructions and results in [12, p. 1759]. The procedure is as follows.

1. Construct a cubic surface S with two skew lines l, m .
2. Calculate the double cover $f : l \rightarrow m$ sending $p \in l$ to the intersection $T_p S \cap m$ of the tangent plane $T_p S$ to p at S with the line m , and construct $g : m \rightarrow l$ analogously.
3. Let $B_f \subset m$ (resp. $B_g \subset l$) be the branch divisor of f (resp. g). Construct morphisms $f' : l \rightarrow \mathbb{P}^1$ (resp. $g' : m \rightarrow \mathbb{P}^1$) ramifying over B_g (resp. B_f).
4. Construct $Q \in m \subset S$ such that $f^{-1}(Q)$ is also a fiber of f' .
5. Construct the ramification locus of the degree 2 projection $S \rightarrow \mathbb{P}(T_Q S)$ from the point $Q \in S$. Then by Proposition 3.1(i) of [12], we obtain a quartic in \mathcal{L}_1 .

The following string of propositions and remarks show how these steps can be implemented without extending the base field (we will take $k = \mathbb{Q}$ throughout).

PROPOSITION 3. *Let k be a field, and suppose that we are given six rational points $p_1, \dots, p_6 \in \mathbb{P}V(k)$ in the projective plane over k . Suppose additionally that this set of points is sufficiently general in the sense that the complete linear system C of cubics passing through them has dimension 4. Construct the Clebsch rational map $c : \mathbb{P}V \rightarrow \mathbb{P}C$, and let $S \subset \mathbb{P}C$ be the Zariski closure of $c(\mathbb{P}V)$. Then S is the vanishing locus of a quaternary cubic form $F \in \text{Sym}^3(C^*)$ over k .*

The rational map c restricts to a birational map between $\mathbb{P}V$ and S . Let $l_0 \subset \mathbb{P}V$ be the rational line containing p_1 and p_2 , and let $m_0 \subset \mathbb{P}V$ be the rational line containing p_1 and p_3 . Let x and x' be two points in l not equal to p_1 or p_2 , and let y and y' be two points in l not equal to p_1 or p_3 . Then the images $c(x)$ and $c(x')$ are well-defined elements of S , and the line l through them is defined over k and included in S . Analogously, one obtains a line m through $c(y)$ and $c(y')$. The lines l and m are skew.

PROOF. This is a standard result from the theory of cubic surfaces, see [6, Section V.4]. \square

This deals with part 1. To perform the calculations in point 2 explicitly, choose coordinates on l by taking two points l_1, l_2 on l and sending $(x : y) \in \mathbb{P}^1$ to $xl_1 + yl_2$, and similarly on m by choosing $m_1, m_2 \in m$. To determine the morphism $f : l \rightarrow m$ explicitly in these coordinates, we choose two equations $M_1 = M_2 = 0$ defining m . Given $p \in l$ with coordinates $(x : y) \in \mathbb{P}^1$, the point $f(p) = T_p S \cap m$ corresponds to the vector space that is the kernel of the matrix whose rows are given by M_1, M_2 and the partial derivatives of F . A generating vector for this space will be a combination of m_1 and m_2 with homogeneous quadratic coefficients $f_1(x, y), f_2(x, y)$ in (x, y) . The morphism f now corresponds to the map $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ given by (f_1, f_2) . Similarly, one determines g . For point 3, we use the following result.

PROPOSITION 4. *Let l be a projective line over k , with homogeneous coordinates x and y , and let D be a k -rational divisor of degree 2 on l . Then the following formulae determine a degree 2 morphism $f' : l \rightarrow \mathbb{P}^1$ over k whose ramification locus equals D .*

- *If D consists of two points $(x_1 : y_1)$ and $(x_2 : y_2)$ that are rational over k , then one can take*

$$f'(x : y) = ((y_1x - x_1y)^2 : (y_2x - x_2y)^2).$$

- *If D is defined by an equation $rx^2 + ty^2 = 0$, then one can take*

$$f'(x : y) = (rx^2 - 2txy - ty^2 : rx^2 + 2txy - ty^2);$$

- *If D is defined by an equation $rx^2 + sxy + ty^2 = 0$ with $s \neq 0$, then one can take*

$$f'(x : y) = (r^2sx^2 + 2r(s^2 - 2rt)xy + (s^3 - 3rst)y^2 : r(rsx^2 + 4rtxy + sty^2)).$$

PROOF. Once the answer is given, the verification is trivial. But let us illustrate how to find these expressions by treating the third case, where the points of D are defined over a proper quadratic extension of k . We use the affine coordinate $t = x/y$ on l . Suppose that using this coordinate, the divisor $D = [d] + [\bar{d}]$ consists of two conjugate points, not summing to zero because we are in case (iii). Then $f' = ((t - d)/(t - \bar{d}))^2$ of above now has the property that $\bar{f}' = 1/f'$. But then one verifies that $(df' + d)/(\bar{d}f' + d)$ is a fractional linear transformation of f' that is stable under conjugation and hence defines a morphism over the ground field. Homogenizing, one obtains the more elegant expression given in the statement of the proposition. \square

We now treat point 4.

PROPOSITION 5. *Let $f, f' : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be two degree 2 morphisms, neither of which can be obtained from the other by postcomposing with an automorphism of \mathbb{P}^1 . Then the q in \mathbb{P}^1 such that the fiber of f over q is also a fiber of f' over a point q' can be obtained as follows.*

Write

$$f(x : y) = (a_1x^2 + b_1xy + c_1y^2 : a_2x^2 + b_2xy + c_2y^2)$$

and

$$f'(x : y) = (a'_1x^2 + b'_1xy + c'_1y^2 : a'_2x^2 + b'_2xy + c'_2y^2).$$

Then $q = (\lambda_1 : \lambda_2)$, where $(\lambda_1, \lambda_2, \lambda'_1, \lambda'_2)$ generates the kernel of the matrix

$$\begin{pmatrix} a_1 & -a_2 & -a'_1 & a'_2 \\ b_1 & -b_2 & -b'_1 & b'_2 \\ c_1 & -c_2 & -c'_1 & c'_2 \end{pmatrix}.$$

PROOF. If we let $q = (\lambda_1 : \lambda_2)$ and $q' = (\lambda'_1 : \lambda'_2)$, then finding q and q' comes down to solving the equation

$$\begin{aligned} & \lambda_2(a_1x^2 + b_1xy + c_1y^2) - \lambda_1(a_2x^2 + b_2xy + c_2y^2) \\ &= \lambda'_2(a'_1x^2 + b'_1xy + c'_1y^2) - \lambda'_1(a'_2x^2 + b'_2xy + c'_2y^2), \end{aligned}$$

which evidently corresponds to the determination of the kernel of the matrix in question. \square

To calculate point 5, we choose an isomorphism $\mathbb{P}C \cong \mathbb{P}^3$ mapping the point Q to $(1 : 0 : 0 : 0)$ and apply the following elementary result.

PROPOSITION 6. *Let $S \subset \mathbb{P}^3$ be a cubic surface containing $Q = (1 : 0 : 0 : 0)$ that is defined by a quaternary cubic form $F \in k[w, x, y, z]$. Let q be the projection $S \rightarrow \mathbb{P}(T_Q S)$ from the point $Q \in S$. Then the ramification locus of q is isomorphic to the quartic curve in \mathbb{P}^2 determined by the vanishing of the discriminant of the quadratic polynomial $F(1, xt, yt, zt)/t$.*

PROOF. Since S is a subscheme of \mathbb{P}^3 , we get an induced coordinatization of $T_Q S$ by sending $(x : y : z)$ to the tangent direction given by the line through the points $(1 : 0 : 0 : 0)$ and $(1 : x : y : z)$. Then $(x : y : z)$ is a ramification point of the projection $S \rightarrow \mathbb{P}(T_Q S)$ if and only if the equation $F(1, xt, yt, zt) = 0$ has a double root outside 0, or in other words if the discriminant of the quadratic polynomial $F(1, xt, yt, zt)/t$ vanishes. This discriminant is a homogeneous quartic form in the variables x, y, z , which defines the plane quartic in \mathcal{L}_1 that we were looking for. \square

A program to generate quartics in \mathcal{L}_1 by using the steps above is available online⁴.

REMARK 1. *We also tried to generate quartics in \mathcal{L}_1 by using Remark 3.4 of [12]. We take cubics S of the form*

$$t^2x + t(ax^2 + 2bxy + 2cxz + dy^2 + 2eyz + fz^2) + g(x, y, z)$$

with g a random degree 3 homogeneous polynomial, such that S is non singular and $e^2 = df$. The last condition ensures that $p = (0 : 0 : 0 : 1)$ belongs to the Hessian H of S and, after checking that p is non singular, we take quartics which are tangent plane sections of H at p . Unfortunately, it seems that these quartics are special in \mathcal{L}_1 , since there are degree 24 relations between their invariants (there is a 27 dimensional space of relations in degree 24 between randomly generated quartics of this form).

Having generated a sufficiently large database⁵ of curves in \mathcal{L}_1 by choosing random 6-tuples $\{p_1, \dots, p_6\}$, we can again proceed as in Section 2. Up to degree 30, all invariants vanishing on the quartics in this databases for \mathcal{L}_1 and \mathcal{L}_2 are multiples of I_{27} . Since the codimension 2 components $\mathcal{L}_1, \mathcal{L}_2, (\mathcal{L}_2)_{\text{red}}$ of $\mathcal{L} \cap \mathcal{D}$ have degree at most $24 \cdot 27, 24 \cdot 30, 24 \cdot 15$, which are all smaller than $(30)^2$, we have the following result.

THEOREM 2. *The subschemes $\mathcal{L}_1, \mathcal{L}_2, (\mathcal{L}_2)_{\text{red}}$ of the projective space of quartic curves $\mathbb{P}\text{Sym}^4(V^*)$ (and hence their images in the coarse moduli space of plane quartic curves) are not complete intersections. In particular, they are not principal hypersurfaces in the discriminant locus \mathcal{D} .*

As there is no degree 24 invariant vanishing on \mathcal{L}_1 , Morley’s putative construction of such an invariant I_{24} in [9, p.282] is incorrect. On the authors’ webpage, a Magma program⁶ is available to check all steps on the way to Theorem 2.

5. OPEN QUESTIONS

The expression L of the Lüroth invariant that we found depends on several arbitrary choices that may explain its

⁴<http://iml.univ-mrs.fr/~ritzenth/programme/luroth/GenerateL1.m>

⁵1024 curves over \mathbb{Q} are available at <http://iml.univ-mrs.fr/~ritzenth/programme/luroth/L1Database.m>

⁶<http://iml.univ-mrs.fr/~ritzenth/programme/luroth/SingularLurothInv.m>

cumbersomeness. First, there is the choice of the basis of invariants. Though some of the Dixmier invariants have geometrical interpretations that are ‘natural’, the same is far from evident for the new Ohno invariants. Secondly, our choice can be modified by any element of the kernel N_1 . Beyond a cancellation of the coefficients of 215 of these monomials that we have already accomplished by simple linear algebra, further minimization of the number of monomials in the expression for L could in theory be achieved by techniques based on coding theory. Still, the parameters seem too large to make this feasible in practice.

The negative answers concerning the existence of degree 24 and 30 invariants in Section 4 exclude the decomposition from [12, p.1764]. The geometry of the situation does not seem to give a clue for the existence of another such decomposition.

An expression in terms of the 15 coefficients of the generic quartic would of course be useful. However, it is not even practically achievable to formally express the fundamental Dixmier-Ohno invariants in this way, since these expressions contain far too many monomials, as is for instance the case for the discriminant I_{27} . A count of weighted monomials in 15 variables for degree 54 invariants leads to a total of 62 422 531 333. Of course only a fraction of these monomials may occur in the final expression of L , but we could not figure out their number, let alone the Newton polytope of L .

Acknowledgments. This note arose from Giorgio Ottaviani’s great lectures at a workshop on invariant theory and projective geometry held in Trento in September 2012, in which the main problems treated in it were mentioned. The authors wish cordially to thank Ottaviani for his further elaborations on the subject, as well as his careful reading of a first draft of this article.

The authors acknowledge support by the grants ANR-09-BLAN-0020-01. The fourth author was additionally supported by the Marie Curie Fellowship IEF-GA-2011-299887.

6. REFERENCES

- [1] W. Barth. Moduli of vector bundles on the projective plane. *Invent. Math.*, 42:63–91, 1977.
- [2] C. Böhnig and H.-C. G. von Bothmer. On the rationality of the moduli space of Lüroth quartics. *Math. Ann.*, 353(4):1273–1281, 2012.
- [3] J. Dixmier. On the projective invariants of quartic plane curves. *Adv. in Math.*, 64:279–304, 1987.
- [4] A.-S. Elsenhans. Explicit computations of invariants of plane quartic curves. To appear in the proceedings of MEGA 2013.
- [5] M. Girard and D. R. Kohel. Classification of genus 3 curves in special strata of the moduli space. Hess, Florian (ed.) et al., Algorithmic number theory. 7th international symposium, ANTS-VII, Berlin, Germany, July 23–28, 2006. Proceedings. Berlin: Springer. Lecture Notes in Computer Science 4076, 346–360, 2006.
- [6] R. Hartshorne. Algebraic Geometry. Springer, 1977.
- [7] J. Hauenstein and F. Sottile. Newton polytopes and witness sets, 2012. Preprint available at arxiv.org/abs/1210.2726.
- [8] J. Le Potier and A. Tikhomirov. Sur le morphisme de Barth. *Annales scientifiques de l’École Normale Supérieure*, 34(4):573–629, 2001.

- [9] F. Morley. On the Lüroth Quartic Curve. *Amer. J. Math.*, 41(4):279–282, 1919.
- [10] T. Ohno. The graded ring of invariants of ternary quartics I, 2005? Unpublished.
- [11] G. Ottaviani and E. Sernesi. On the hypersurface of Lüroth quartics. *Michigan Math. J.*, 59(2):365–394, 2010.
- [12] G. Ottaviani and E. Sernesi. On singular Lüroth quartics. *Sci. China Math.*, 54(8):1757–1766, 2011.
- [13] G. Ottaviani. A computational approach to Lüroth quartics, 2012. *Rendiconti del Circolo Matematico di Palermo* (2)62, no.1, 165–177, 2013.
- [14] A. Popolitov and S. Shakirov. On undulation invariants of plane curves, 2012. Preprint available at <http://arxiv.org/abs/1208.5775>.
- [15] T. Shioda. On the graded ring of invariants of binary octavics. *American J. of Math.*, 89(4):1022–1046, 1967.